antibodies .- online.com





anti-ZFAND5 antibody (AA 121-213)



| () | 1/0 | r\ /1 | 014 | |
|-----|-----|-------|-----|---|
| () | ve | I V I | -v | V |
| | | | | |

| Quantity: | 100 μL | |
|----------------------|--|--|
| Target: | ZFAND5 | |
| Binding Specificity: | AA 121-213 | |
| Reactivity: | Human | |
| Host: | Rabbit | |
| Clonality: | Polyclonal | |
| Conjugate: | This ZFAND5 antibody is un-conjugated | |
| Application: | cation: Immunofluorescence (Cultured Cells) (IF (cc)), Immunofluorescence (Paraffin-embedded Sections) (IF (p)), ELISA, Immunohistochemistry (Paraffin-embedded Sections) (IHC (p)), Immunohistochemistry (Frozen Sections) (IHC (fro)), Immunocytochemistry (ICC) | |

Product Details

| Immunogen: | KLH conjugated synthetic peptide derived from human ZFAND5 |
|-----------------------|--|
| Isotype: | IgG |
| Predicted Reactivity: | Human,Mouse,Rat,Cow,Sheep,Pig,Chicken,Rabbit |
| Purification: | Purified by Protein A. |
| Target Details | |

| Target: | ZFAND5 |
|-------------------|--------------------------|
| Alternative Name: | ZFAND5 (ZFAND5 Products) |

Target Details

| 3 | | | |
|---------------------|--|--|--|
| Background: | Synonyms: ZA20D2, ZNF216, ZFAND5A, AN1-type zinc finger protein 5, Zinc finger A20 domain- | | |
| | containing protein 2, Zinc finger protein 216, ZFAND5 | | |
| | Background: Involved in protein degradation via the ubiquitin-proteasome system. May act by | | |
| | anchoring ubiquitinated proteins to the proteasome. Plays a role in ubiquitin-mediated protein | | |
| | degradation during muscle atrophy. Plays a role in the regulation of NF-kappa-B activation and | | |
| | apoptosis. Inhibits NF-kappa-B activation triggered by overexpression of RIPK1 and TRAF6 but | | |
| | not of RELA. Inhibits also tumor necrosis factor (TNF), IL-1 and TLR4-induced NF-kappa-B | | |
| | activation in a dose-dependent manner. Overexpression sensitizes cells to TNF-induced | | |
| | apoptosis. Is a potent inhibitory factor for osteoclast differentiation. | | |
| Gene ID: | 7763 | | |
| UniProt: | 076080 | | |
| Pathways: | Platelet-derived growth Factor Receptor Signaling | | |
| Application Details | | | |
| Application Notes: | ELISA 1:500-1000 | | |
| | IHC-P 1:200-400 | | |
| | IHC-F 1:100-500 | | |
| | IF(IHC-P) 1:50-200 | | |
| | IF(IHC-F) 1:50-200 | | |
| | IF(ICC) 1:50-200 | | |
| | ICC 1:100-500 | | |
| Restrictions: | For Research Use only | | |
| Handling | | | |
| Format: | Liquid | | |
| Concentration: | 1 μg/μL | | |
| Buffer: | 0.01M TBS(pH 7.4) with 1 % BSA, 0.02 % Proclin300 and 50 % Glycerol. | | |
| Preservative: | ProClin | | |
| Precaution of Use: | This product contains ProClin: a POISONOUS AND HAZARDOUS SUBSTANCE, which should be handled by trained staff only. | | |
| Storage: | 4 °C,-20 °C | | |
| | | | |

Handling

| Storage Comment: | Shipped at 4°C. Store at -20°C for one year. Avoid repeated freeze/thaw cycles. |
|------------------|---|
| Expiry Date: | 12 months |