# antibodies .- online.com







## anti-GMEB1 antibody (AA 301-400)



$\sim$			
	N/P	r\/I	i⊢₩

Quantity:	100 μL
Target:	GMEB1
Binding Specificity:	AA 301-400
Reactivity:	Human
Host:	Rabbit
Clonality:	Polyclonal
Conjugate:	This GMEB1 antibody is un-conjugated
Application:	Western Blotting (WB), ELISA, Immunofluorescence (Cultured Cells) (IF (cc)), Immunofluorescence (Paraffin-embedded Sections) (IF (p)), Immunohistochemistry (Paraffin-embedded Sections) (IHC (p)), Immunocytochemistry (ICC), Immunohistochemistry (Frozen Sections) (IHC (fro))

#### **Product Details**

Immunogen:	KLH conjugated synthetic peptide derived from human GMEB1
Isotype:	IgG
Predicted Reactivity:	Human,Mouse,Rat,Cow,Sheep,Horse,Rabbit
Purification:	Purified by Protein A.

## **Target Details**

Target:	GMEB1
Alternative Name:	GMEB1 (GMEB1 Products)

#### Target Details

Background:
-------------

Synonyms: DNA binding protein p96P, DNA-binding protein p96P, Glucocorticoid modulatory element binding protein 1, Glucocorticoid modulatory element-binding protein 1, GMEB 1, GMEB-1, Gmeb1, GMEB1\_HUMAN, P96P, Parvovirus initiation factor p96, P 96, P p96, P96. Background: GMEB-1 is a 573 amino acid protein that contains one SAND domain and is a member of the KDWK family of combinatorial transcription modulators. Localized to both the cytoplasm and the nucleus, GMEB-1 forms a heterodimer with GMEB-2 (Glucocorticoid modulatory element-binding protein 2) and, once associated with GMEB-2, plays a key role in parvovirus DNA replication. In addition, GMEB-1 functions alone as a trans-acting factor that, by binding to glucocorticoid modulatory elements (GMEs) in TAT (tyrosine aminotransferase) promoters, increases intracellular sensitivity to glucocorticoid concentrations. GMEB-1 also interacts with initiator procaspases and, via this interaction, can inhibit caspase-induced apoptosis. Due to alternative splicing events, GMEB-1 is expressed as two isoforms.

Gene ID:

10691

#### **Application Details**

Application Notes:	

WB 1:300-5000

ELISA 1:500-1000

IHC-P 1:200-400

IHC-F 1:100-500

IF(IHC-P) 1:50-200

IF(IHC-F) 1:50-200

IF(ICC) 1:50-200

ICC 1:100-500

Restrictions:

For Research Use only

### Handling

Format:	Liquid	
Concentration:	1 μg/μL	
Buffer:	0.01M TBS( pH 7.4) with 1 % BSA, 0.02 % Proclin300 and 50 % Glycerol.	
Preservative:	ProClin	
Precaution of Use:	This product contains ProClin: a POISONOUS AND HAZARDOUS SUBSTANCE, which should be handled by trained staff only.	

## Handling

Storage:	4 °C,-20 °C
Storage Comment:	Shipped at 4°C. Store at -20°C for one year. Avoid repeated freeze/thaw cycles.
Expiry Date:	12 months