antibodies.com

# Datasheet for ABIN1714418 anti-ATPase Inhibitory Factor 1 antibody (AA 26-106)



Overview

Quantity:	100 μL
Target:	ATPase Inhibitory Factor 1 (ATPIF1)
Binding Specificity:	AA 26-106
Reactivity:	Rat
Host:	Rabbit
Clonality:	Polyclonal
Conjugate:	This ATPase Inhibitory Factor 1 antibody is un-conjugated
Application:	Western Blotting (WB), ELISA

### Product Details

Immunogen:	KLH conjugated synthetic peptide derived from human ATPIF1/ATPase Inhibitory Factor 1
lsotype:	lgG
Cross-Reactivity:	Human
Predicted Reactivity:	Mouse,Rat,Dog,Horse,Rabbit
Purification:	Purified by Protein A.

## Target Details

Target:	ATPase Inhibitory Factor 1 (ATPIF1)
Alternative Name:	ATP1/ATPase Inhibitory Factor 1 (ATPIF1 Products)
Background:	Synonyms: AT1_HUMAN, ATP synthase inhibitor protein, ATPase inhibitor, ATPase inhibitor

Order at www.antibodies-online.com | www.antikoerper-online.de | www.anticorps-enligne.fr | www.antibodies-online.cn International: +49 (0)241 95 163 153 | USA & Canada: +1 877 302 8632 | support@antibodies-online.com Page 1/3 | Product datasheet for ABIN1714418 | 03/07/2024 | Copyright antibodies-online. All rights reserved. mitochondrial, ATPase inhibitor protein, ATPase inhibitory factor 1, ATPI, ATPI, ATPI, ATPIP, 1, 1, Inhibitor of F1Fo-ATPase, IP, MGC1167, MGC8898, mitochondrial. Background: Mitochondrial ATP synthases (ATPases) transduce the energy contained in membrane electrochemical proton gradients into the energy required for synthesis of highenergy phosphate bonds. ATPases contain two linked complexes: F1, the hydrophilic catalytic core, and F0, the membrane-embedded protein channel. F1 consists of three Alpha chains and three Beta chains, which are weakly homologous, as well as one Gamma chain, one Delta chain and one Gamma chain. F0 consists of three subunits: a, b and c. A mitochondrial F1-ATPase inhibitor protein, ATPIF1 (ATPase inhibitory factor 1), also known as IP, IF1, ATPI or ATPIP (ATPase inhibitor protein), binds to the C-terminal region of a Beta subunit of the F1-ATPase at low pH values and, via interference of the Beta and Gamma subunit interaction, ATPIF1 regulates the activity of the F1F0-ATPase. This reversible ATPIF1 binding to F1F0-ATPase also occurs on the surface of endothelial cells.

Gene ID:

93974

#### Application Details

Application Notes:	ELISA 1:500-1000
	IHC-P 1:200-400
	IHC-F 1:100-500
	IF(IHC-P) 1:50-200
	IF(IHC-F) 1:50-200
	IF(ICC) 1:50-200
	ICC 1:100-500
Restrictions:	For Research Use only

#### Handling

Format:	Liquid
Concentration:	1 μg/μL
Buffer:	0.01M TBS( pH 7.4) with 1 % BSA, 0.02 % Proclin300 and 50 % Glycerol.
Preservative:	ProClin
Precaution of Use:	This product contains ProClin: a POISONOUS AND HAZARDOUS SUBSTANCE, which should be handled by trained staff only.

Order at www.antibodies-online.com | www.antikoerper-online.de | www.anticorps-enligne.fr | www.antibodies-online.cn International: +49 (0)241 95 163 153 | USA & Canada: +1 877 302 8632 | support@antibodies-online.com Page 2/3 | Product datasheet for ABIN1714418 | 03/07/2024 | Copyright antibodies-online. All rights reserved.

Handling	
Storage:	4 °C,-20 °C
Storage Comment:	Shipped at 4°C. Store at -20°C for one year. Avoid repeated freeze/thaw cycles.
Expiry Date:	12 months