.-online.com antibodies

Datasheet for ABIN1714441 anti-EXTL3 antibody (AA 351-450)



Overview

Quantity:	100 µL
Target:	EXTL3
Binding Specificity:	AA 351-450
Reactivity:	Rat
Host:	Rabbit
Clonality:	Polyclonal
Conjugate:	This EXTL3 antibody is un-conjugated
Application:	Western Blotting (WB), ELISA, Immunohistochemistry (Paraffin-embedded Sections) (IHC (p)), Immunofluorescence (Cultured Cells) (IF (cc)), Immunofluorescence (Paraffin-embedded Sections) (IF (p)), Immunohistochemistry (Frozen Sections) (IHC (fro)), Immunocytochemistry (ICC)

Product Details

Immunogen:	KLH conjugated synthetic peptide derived from human EXTL3
Isotype:	lgG
Cross-Reactivity:	Rat
Predicted Reactivity:	Human,Mouse,Dog,Horse
Purification:	Purified by Protein A.
Target Details	

Target:

EXTL3

Order at www.antibodies-online.com | www.antikoerper-online.de | www.anticorps-enligne.fr | www.antibodies-online.cn International: +49 (0)241 95 163 153 | USA & Canada: +1 877 302 8632 | support@antibodies-online.com Page 1/3 | Product datasheet for ABIN1714441 | 03/08/2024 | Copyright antibodies-online. All rights reserved.

Target Details	
Alternative Name:	EXTL3 (EXTL3 Products)
Background:	Synonyms: botv, DKFZp686C2342, Exostoses multiple-like 3, Exostoses-like 3, Exostosin-like 3
	EXT-related protein 1, EXTL1L, EXTL3, EXTL3_HUMAN, EXTR1, Glucuronyl-galactosyl-
	proteoglycan 4-alpha-N-acetylglucosaminyltransferas, Glucuronyl-galactosyl-proteoglycan 4-
	alpha-N-acetylglucosaminyltransferase, Hereditary multiple exostoses gene isolog, KIAA0519,
	Multiple exostosis-like protein 3, Putative tumor suppressor protein EXTL3, REG, Reg receptor, REGR, RPR.
	Background: EXTL3 is a member of the EXT (hereditary multiple exostosin) gene family of
	tumor suppressors encoding glycosyltransferases involved in heparan sulfate (HS)
	biosynthesis. Within this family, the C-terminus is conserved between all members from C.
	elegans to vertebrates. EXTL3 is a ubiquitously expressed, developmentally regulated, single-
	pass type II membrane protein that localizes to the endoplasmic reticulum membrane. EXTL3
	adds N-acetylglucosamine (GlcNAc) to the polysaccharide-protein linkage region and to the
	growing HS chain suggesting that it plays a role in both the initiation and elongation of HS
	chains. In addition, EXTL3 may act as a Reg receptor, binding Reg via its N-terminus.
Gene ID:	2137
Pathways:	Glycosaminoglycan Metabolic Process, ER-Nucleus Signaling
Application Details	
Application Notes:	WB 1:300-5000
	ELISA 1:500-1000
	IHC-P 1:200-400
	IHC-F 1:100-500
	IF(IHC-P) 1:50-200
	IF(IHC-F) 1:50-200
	IF(ICC) 1:50-200
	ICC 1:100-500
Restrictions:	For Research Use only
Handling	
Format:	Liquid

Concentration:	1 μg/μL
Buffer:	0.01M TBS(pH 7.4) with 1 % BSA, 0.02 % Proclin300 and 50 % Glycerol.
	dies-online.com www.antikoerper-online.de www.anticorps-enligne.fr www.antibodies-online.cn +49 (0)241 95 163 153 USA & Canada: +1 877 302 8632 support@antibodies-online.com

Page 2/3 | Product datasheet for ABIN1714441 | 03/08/2024 | Copyright antibodies-online. All rights reserved.

Handling

Preservative:	ProClin
Precaution of Use:	This product contains ProClin: a POISONOUS AND HAZARDOUS SUBSTANCE, which should be handled by trained staff only.
Storage:	4 °C,-20 °C
Storage Comment:	Shipped at 4°C. Store at -20°C for one year. Avoid repeated freeze/thaw cycles.
Expiry Date:	12 months