

## Datasheet for ABIN1714477

# anti-PIK3 gamma antibody (AA 521-620)



#### Overview

Quantity:	100 μL
Target:	PIK3 gamma (PIK3CG)
Binding Specificity:	AA 521-620
Reactivity:	Human
Host:	Rabbit
Clonality:	Polyclonal
Conjugate:	This PIK3 gamma antibody is un-conjugated
Application:	Western Blotting (WB), ELISA, Immunofluorescence (Cultured Cells) (IF (cc)), Immunofluorescence (Paraffin-embedded Sections) (IF (p)), Immunohistochemistry (Paraffin-embedded Sections) (IHC (p)), Flow Cytometry (FACS), Immunohistochemistry (Frozen Sections) (IHC (fro)), Immunocytochemistry (ICC)

#### **Product Details**

Immunogen:	KLH conjugated synthetic peptide derived from human PIK3 gamma
Isotype:	IgG
Cross-Reactivity:	Human
Predicted Reactivity:	Mouse,Rat,Dog,Rabbit
Purification:	Purified by Protein A.

## **Target Details**

Target: PIK3 gamma (PIK3CG)

## Target Details

Abstract:	PIK3CG Products
Background:	Synonyms: PK3CG_HUMAN, PI 3 Kinase catalytic subunit gamma, Phosphatidylinositol 4,5-
	bisphosphate 3-kinase catalytic subunit gamma isoform, PI3-kinase subunit gamma, PI3K-
	gamma, PI3Kgamma,
	Background: This gene encodes a protein that belongs to the pi3/pi4-kinase family of proteins.
	The gene product is an enzyme that phosphorylates phosphoinositides on the 3-hydroxyl group
	of the inositol ring. It is an important modulator of extracellular signals, including those elicited
	by E-cadherin-mediated cell-cell adhesion, which plays an important role in maintenance of the
	structural and functional integrity of epithelia. In addition to its role in promoting assembly of
	adherens junctions, the protein is thought to play a pivotal role in the regulation of cytotoxicity
	in NK cells. The gene is located in a commonly deleted segment of chromosome 7 previously
	identified in myeloid leukemias. [provided by RefSeq, Jul 2008].
Gene ID:	5294
UniProt:	P48736
Pathways:	PI3K-Akt Signaling, RTK Signaling, AMPK Signaling, TLR Signaling, Inositol Metabolic Process,
	Hepatitis C, VEGF Signaling

# **Application Details**

Application Notes:	WB 1:300-5000
	ELISA 1:500-1000
	FCM 1:20-100
	IHC-P 1:200-400
	IHC-F 1:100-500
	IF(IHC-P) 1:50-200
	IF(IHC-F) 1:50-200
	IF(ICC) 1:50-200
	ICC 1:100-500
Restrictions:	For Research Use only
Handling	
Format:	Liquid
Concentration:	1 μg/μL

## Handling

Buffer:	0.01M TBS(pH 7.4) with 1 % BSA, 0.02 % Proclin300 and 50 % Glycerol.
Preservative:	ProClin
Precaution of Use:	This product contains ProClin: a POISONOUS AND HAZARDOUS SUBSTANCE, which should be handled by trained staff only.
Storage:	4 °C,-20 °C
Storage Comment:	Shipped at 4°C. Store at -20°C for one year. Avoid repeated freeze/thaw cycles.
Expiry Date:	12 months