antibodies -online.com







Overview

| Quantity: | 100 μL |
|----------------------|---|
| Target: | BANF1 |
| Binding Specificity: | AA 21-89 |
| Reactivity: | Human |
| Host: | Rabbit |
| Clonality: | Polyclonal |
| Conjugate: | This BANF1 antibody is un-conjugated |
| Application: | Western Blotting (WB), ELISA, Immunofluorescence (Cultured Cells) (IF (cc)), Immunofluorescence (Paraffin-embedded Sections) (IF (p)), Immunohistochemistry (Paraffin-embedded Sections) (IHC (p)), Immunocytochemistry (ICC), Immunohistochemistry (Frozen Sections) (IHC (fro)) |

Product Details

| Immunogen: | KLH conjugated synthetic peptide derived from human BANF1/BAF |
|-----------------------|---|
| Isotype: | IgG |
| Predicted Reactivity: | Human,Mouse,Rat,Dog,Cow |
| Purification: | Purified by Protein A. |

Target Details

| Target: | BANF1 |
|-------------------|----------------------------|
| Alternative Name: | BANF1/BAF (BANF1 Products) |

Target Details

| Background: |
|-------------|
|-------------|

Synonyms: BAF, BAF_HUMAN, BANF 1, BANF1, Barrier to autointegration factor 1, Barrier to autointegration factor, Barrier-to-autointegration factor, BCRG 1, BCRG1, BCRP 1, BCRP1, Breakpoint cluster region protein 1, D14S1460, MGC111161.

Background: Barrier-to-autointegration factor (BAF) binds non-specifically to double stranded DNA, possibly to play a role in tissue- or cell type-specific gene expression by interacting with different homeodomain transcription factors. BAF compresses chromatin structure and interacts with the LEM domain of nuclear proteins to play a crucial role in membrane recruitment and chromatin decondensation during nuclear assembly. Additionally, retroviruses like HIV-1 incorporate BAF from host cells into preintegration complexes (PICs) to prevent autointegration of retroviral DNA and thereby promote integration of retroviral DNA into the host chromosome.

Gene ID:

8815

Application Details

| Application Notes: | WB 1:300-5000 |
|--------------------|---------------|
| | |

ELISA 1:500-1000

IHC-P 1:200-400

IHC-F 1:100-500

IF(IHC-P) 1:50-200

IF(IHC-F) 1:50-200

IF(ICC) 1:50-200

ICC 1:100-500

Restrictions:

For Research Use only

Handling

| Format: | Liquid |
|--------------------|--|
| Concentration: | 1 μg/μL |
| Buffer: | 0.01M TBS(pH 7.4) with 1 % BSA, 0.02 % Proclin300 and 50 % Glycerol. |
| Preservative: | ProClin |
| Precaution of Use: | This product contains ProClin: a POISONOUS AND HAZARDOUS SUBSTANCE, which should be handled by trained staff only. |
| Storage: | 4 °C,-20 °C |

Handling

| Storage Comment: | Shipped at 4°C. Store at -20°C for one year. Avoid repeated freeze/thaw cycles. |
|------------------|---|
| Expiry Date: | 12 months |