antibodies .- online.com





Entrates and the second

Datasheet for ABIN1714499

anti-C1orf87 antibody (AA 401-500)

Overview Quantity:

100 μL

Target: C1orf87

Binding Specificity: AA 401-500

Reactivity: Human

Host: Rabbit

Clonality: Polyclonal

Conjugate: This C1orf87 antibody is un-conjugated

Application: ELISA, Immunofluorescence (Cultured Cells) (IF (cc)), Immunofluorescence (Paraffin-

embedded Sections) (IF (p)), Western Blotting (WB), Immunohistochemistry (Frozen Sections)

(IHC (fro)), Immunohistochemistry (Paraffin-embedded Sections) (IHC (p)),

Immunocytochemistry (ICC)

Product Details

Immunogen:	KLH conjugated synthetic peptide derived from human C1orf87
------------	---

Isotype: IgG

Predicted Reactivity: Human, Mouse, Rat, Dog, Cow, Sheep

Purification: Purified by Protein A.

Target Details

Target: C1orf87

Alternative Name: C1orf87 (C1orf87 Products)

Target Details

Background:

Synonyms: chromosome 1 open reading frame 87, FLJ44073, LOC127795, MGC34837, Uncharacterized protein C1orf87, CA087_HUMAN.

Background: Chromosome 1 is the largest human chromosome spanning about 260 million base pairs and making up 8 % of the human genome. There are about 3,000 genes on chromosome 1, and considering the great number of genes there are also a large number of diseases associated with chromosome 1. Notably, the rare aging disease Hutchinson-Gilford progeria is associated with the LMNA gene which encodes lamin A. When defective, the LMNA gene product can build up in the nucleus and cause characteristic nuclear blebs. The mechanism of rapidly enhanced aging is unclear and is a topic of continuing exploration. The MUTYH gene is located on chromosome 1 and is partially responsible for familial adenomatous polyposis. Stickler syndrome, Parkinsons, Gaucher disease and Usher syndrome are also associated with chromosome 1. A breakpoint has been identified in 1q which disrupts the DISC1 gene and is linked to schizophrenia. Aberrations in chromosome 1 are found in a variety of cancers including head and neck cancer, malignant melanoma and multiple myeloma. The C1orf87 gene product has been provisionally designated C1orf87 pending further characterization. There are three isoforms of C1orf87 that are produced as a result of alternative splicing events.

Gene ID:

127795

Application Details

Application Notes: WB 1:300-5000

ELISA 1:500-1000

IHC-P 1:200-400

IHC-F 1:100-500

IF(IHC-P) 1:50-200

IF(IHC-F) 1:50-200

IF(ICC) 1:50-200

ICC 1:100-500

Restrictions:

For Research Use only

Handling

Format: Liquid

Concentration: $1 \mu g/\mu L$

Handling

Buffer:	0.01M TBS(pH 7.4) with 1 % BSA, 0.02 % Proclin300 and 50 % Glycerol.
Preservative:	ProClin
Precaution of Use:	This product contains ProClin: a POISONOUS AND HAZARDOUS SUBSTANCE, which should be handled by trained staff only.
Storage:	4 °C,-20 °C
Storage Comment:	Shipped at 4°C. Store at -20°C for one year. Avoid repeated freeze/thaw cycles.
Expiry Date:	12 months