antibodies -online.com





Datasheet for ABIN1714608 anti-CACFD1 antibody (AA 41-130)

Go to Product page

Overview

Quantity:	100 μL
Target:	CACFD1
Binding Specificity:	AA 41-130
Reactivity:	Human
Host:	Rabbit
Clonality:	Polyclonal
Conjugate:	This CACFD1 antibody is un-conjugated
Application:	Western Blotting (WB), ELISA, Immunofluorescence (Cultured Cells) (IF (cc)), Immunofluorescence (Paraffin-embedded Sections) (IF (p)), Immunohistochemistry (Paraffin-embedded Sections) (IHC (p)), Immunohistochemistry (Frozen Sections) (IHC (fro)), Immunocytochemistry (ICC)

Product Details

Immunogen:	KLH conjugated synthetic peptide derived from human C9orf7
Isotype:	IgG
Predicted Reactivity:	Human,Mouse,Rat,Dog,Pig,Rabbit
Purification:	Purified by Protein A.

Target Details

Target:	CACFD1
Alternative Name:	C9orf7 (CACFD1 Products)

Target Details

Bac	kar	ound:

Synonyms: C9orf7, Calcium channel flower homolog, Chromosome 9 open reading frame 7, D9S2135, FLOWR_HUMAN, RP13-100B2.5.

Background: Chromosome 9 consists of about 145 million bases and 4 % of the human genome and encodes nearly 900 genes. Considered to play a role in gender determination, deletion of the distal portion of 9p can lead to development of male to female sex reversal, the phenotype of a female with a male X,Y genotype. Hereditary hemorrhagic telangiectasia, which is characterized by harmful vascular defects, is associated with the chromosome 9 gene encoding endoglin protein, ENG. Familial dysautonomia is also associated with chromosome 9 though through the gene IKBKAP. Notably, chromosome 9 encompasses the largest interferon family gene cluster. Chromosome 9 is partnered with chromosome 22 in the translocation leading to the aberrant production of BCR-ABL fusion protein often found in leukemias. The C9orf7 gene product has been provisionally designated C9orf7 pending further characterization.

Gene ID:

11094

Application Details

Application Notes:

WB 1:300-5000

ELISA 1:500-1000

IHC-P 1:200-400

IHC-F 1:100-500

IF(IHC-P) 1:50-200

IF(IHC-F) 1:50-200

IF(ICC) 1:50-200

ICC 1:100-500

Restrictions:

For Research Use only

Handling

Format:	Liquid
Concentration:	1 μg/μL
Buffer:	0.01M TBS(pH 7.4) with 1 % BSA, 0.02 % Proclin300 and 50 % Glycerol.
Preservative:	ProClin
Precaution of Use:	This product contains ProClin: a POISONOUS AND HAZARDOUS SUBSTANCE, which should be handled by trained staff only.

Handling

Storage:	4 °C,-20 °C	
Storage Comment:	Shipped at 4°C. Store at -20°C for one year. Avoid repeated freeze/thaw cycles.	
Expiry Date:	12 months	