antibodies -online.com





Datasheet for ABIN1714615

anti-FOXN2 antibody (AA 51-150)



Overview

Quantity:	100 μL
Target:	FOXN2
Binding Specificity:	AA 51-150
Reactivity:	Mouse
Host:	Rabbit
Clonality:	Polyclonal
Conjugate:	This FOXN2 antibody is un-conjugated
Application:	Western Blotting (WB), ELISA, Immunofluorescence (Paraffin-embedded Sections) (IF (p)), Immunofluorescence (Cultured Cells) (IF (cc)), Immunohistochemistry (Paraffin-embedded Sections) (IHC (p)), Immunohistochemistry (Frozen Sections) (IHC (fro)), Immunocytochemistry (ICC)

Product Details

Immunogen:	KLH conjugated synthetic peptide derived from human FOXN2
Isotype:	IgG
Cross-Reactivity:	Mouse
Predicted Reactivity:	Human,Rat,Dog,Pig,Rabbit
Purification:	Purified by Protein A.

Target Details

Target: FOXN2

Target Details

Alternative Name:	FOXN2 (FOXN2 Products)
Background:	Synonyms: 3230402J05Rik, 6030465J18Rik, Fkh19, Forkhead box N2, Forkhead box protein
	N2, Foxn1, foxn2, FOXN2_HUMAN, HTLF, Human T-cell leukemia virus enhancer factor.
	Background: The forkhead domain-containing gene family (Fox) comprises over 20 members in
	mammals and is defined by a conserved 110 amino-acid motif containing a winged helix
	structure DNA-binding domain. The members of this gene family have been implicated as key
	regulators of embryogenesis, cell cycling, cell lineage restriction and cancer. As such, FOXN2
	contains a domain with homology to the forkhead DNA binding domain. FOXN2, or Human T-
	cell leukemia virus enhancer factor, is a 341 amino acid protein mapping to human gene
	FOXN2, which has been localized to human chromosome 2p16-p22. This protein, encoded by a
	1239-bp cDNA isolated from the Jurkat cDNA library, is capable of binding to a region of the
	human T-cell leukemia virus long terminal repeat (HTLV-I LTR) located between amino acids
	155 and 117. This purine-rich region is important in the regulation of gene expression by the Et
	family of transcription factors. FOXN2 is a unique cellular gene that may function in the
	transcriptional regulation of HTLV-I LTR.
Gene ID:	3344
Application Details	
Application Notes:	WB 1:300-5000
	ELISA 1:500-1000
	IHC-P 1:200-400
	IHC-F 1:100-500
	IF(IHC-P) 1:50-200
	IF(IHC-F) 1:50-200
	IF(ICC) 1:50-200
	ICC 1:100-500
Restrictions:	For Research Use only
Handling	
Format:	Liquid
Concentration:	1 μg/μL
Buffer:	0.01M TBS(pH 7.4) with 1 % BSA, 0.02 % Proclin300 and 50 % Glycerol.
Preservative:	ProClin

Handling

Precaution of Use:	This product contains ProClin: a POISONOUS AND HAZARDOUS SUBSTANCE, which should be handled by trained staff only.
Storage:	4 °C,-20 °C
Storage Comment:	Shipped at 4°C. Store at -20°C for one year. Avoid repeated freeze/thaw cycles.
Expiry Date:	12 months