

[Go to Product page](#)

Datasheet for ABIN1714653 anti-NAPA antibody (AA 201-295)

Overview

Quantity:	100 µL
Target:	NAPA
Binding Specificity:	AA 201-295
Reactivity:	Human
Host:	Rabbit
Clonality:	Polyclonal
Conjugate:	This NAPA antibody is un-conjugated
Application:	ELISA, Western Blotting (WB), Immunofluorescence (Cultured Cells) (IF (cc)), Immunofluorescence (Paraffin-embedded Sections) (IF (p)), Immunocytochemistry (ICC), Immunohistochemistry (Paraffin-embedded Sections) (IHC (p)), Immunohistochemistry (Frozen Sections) (IHC (fro))

Product Details

Immunogen:	KLH conjugated synthetic peptide derived from human Alpha SNAP
Isotype:	IgG
Predicted Reactivity:	Human, Mouse, Rat, Dog, Cow, Pig
Purification:	Purified by Protein A.

Target Details

Target:	NAPA
Alternative Name:	Alpha SNAP (NAPA Products)

Target Details

Background:	<p>Synonyms: Alpha soluble NSF attachment protein, Alpha-soluble NSF attachment protein, N ethylmaleimide sensitive factor attachment protein alpha, N-ethylmaleimide-sensitive factor attachment protein alpha, NAPA, SNAA_HUMAN, SNAP alpha, SNAP alpha, SNAP-alpha, SNAPA.</p> <p>Background: Syntaxins were originally thought to be docking proteins, but have more recently been categorized as anchoring proteins that anchor themselves to the cytoplasmic surfaces of cellular membranes. Syntaxins have been shown to bind to various proteins involved in exocytosis, including VAMPs (vesicle-associated membrane proteins), NSF (N-ethylmaleimide-sensitive factor), SNAP 25 (synaptosomal-associated protein of 25 kDa), SNAPs (soluble NSF attachment proteins) and synaptotagmin. VAMPs (also designated synaptobrevins), including VAMP-1 and VAMP-2, and synaptotagmin, a protein that may function as an inhibitor of exocytosis, are vesicular proteins.</p>
Gene ID:	8775
Pathways:	Synaptic Vesicle Exocytosis , Asymmetric Protein Localization

Application Details

Application Notes:	WB 1:300-5000 ELISA 1:500-1000 IHC-P 1:200-400 IHC-F 1:100-500 IF(IHC-P) 1:50-200 IF(IHC-F) 1:50-200 IF(ICC) 1:50-200 ICC 1:100-500
--------------------	----------------------------------------------------------------------------------------------------------------------------------------------------------

Restrictions:	For Research Use only
---------------	-----------------------

Handling

Format:	Liquid
Concentration:	1 µg/µL
Buffer:	0.01M TBS(pH 7.4) with 1 % BSA, 0.02 % Proclin300 and 50 % Glycerol.
Preservative:	ProClin
Precaution of Use:	This product contains ProClin: a POISONOUS AND HAZARDOUS SUBSTANCE, which should be

Handling

	handled by trained staff only.
Storage:	4 °C,-20 °C
Storage Comment:	Shipped at 4°C. Store at -20°C for one year. Avoid repeated freeze/thaw cycles.
Expiry Date:	12 months