antibodies - online.com







anti-TYROBP antibody (AA 31-113)



/ N	1	-	K	/1	01	A /
	1//	\vdash	1 \	/ I	e	\/\/
\sim	v	\sim	1 V		\sim	v v

Quantity:	100 μL
Target:	TYROBP
Binding Specificity:	AA 31-113
Reactivity:	Mouse
Host:	Rabbit
Clonality:	Polyclonal
Conjugate:	This TYROBP antibody is un-conjugated
Application:	Western Blotting (WB), Flow Cytometry (FACS), ELISA, Immunofluorescence (Cultured Cells) (IF
	(cc)), Immunofluorescence (Paraffin-embedded Sections) (IF (p)), Immunohistochemistry
	(Paraffin-embedded Sections) (IHC (p)), Immunohistochemistry (Frozen Sections) (IHC (fro))

Product Details

Immunogen:	KLH conjugated synthetic peptide derived from human DAP12
Isotype:	IgG
Cross-Reactivity:	Mouse
Predicted Reactivity:	Human,Rat,Dog,Cow,Sheep,Pig,Horse
Purification:	Purified by Protein A.

Target Details

rarget: rykobp

Target Details

Alternative Name:	DAP12 (TYROBP Products)		
Background:	Synonyms: DAP 12, DAP12, DNAX activation protein 12, DNAX-activation protein 12, KAR-		
	associated protein, KARAP, Killer activating receptor associated protein, Killer-activating		
	receptor-associated protein, PLOSL, TYOBP_HUMAN, TYRO protein tyrosine kinase binding		
	protein, TYRO protein tyrosine kinase-binding protein, TYROBP.		
	Background: This gene encodes a transmembrane signaling polypeptide which contains an		
	immunoreceptor tyrosine-based activation motif (ITAM) in its cytoplasmic domain. The		
	encoded protein may associate with the killer-cell inhibitory receptor (KIR) family of membrane		
	glycoproteins and may act as an activating signal transduction element. This protein may bind		
	zeta-chain (TCR) associated protein kinase 70 kDa (ZAP-70) and spleen tyrosine kinase (SYK)		
	and play a role in signal transduction, bone modeling, brain myelination, and inflammation.		
	Mutations within this gene have been associated with polycystic lipomembranous		
	osteodysplasia with sclerosing leukoencephalopathy (PLOSL), also known as Nasu-Hakola		
	disease. Its putative receptor, triggering receptor expressed on myeloid cells 2 (TREM2), also		
	causes PLOSL. Multiple alternative transcript variants encoding distinct isoforms have been		
	identified for this gene. [provided by RefSeq, Mar 2010]		
Gene ID:	7305		
UniProt:	043914		
Application Details			
Application Notes:	WB 1:300-5000		
	ELISA 1:500-1000		
	FCM 1:20-100		
	IHC-P 1:200-400		
	IHC-F 1:100-500		
	IF(IHC-P) 1:50-200		
	IF(IHC-F) 1:50-200		
	IF(ICC) 1:50-200		
Restrictions:	For Research Use only		
Handling			
Format:	Liquid		
Concentration:	1 μg/μL		

Handling

Buffer:	0.01M TBS(pH 7.4) with 1 % BSA, 0.02 % Proclin300 and 50 % Glycerol.
Preservative:	ProClin
Precaution of Use:	This product contains ProClin: a POISONOUS AND HAZARDOUS SUBSTANCE, which should be handled by trained staff only.
Storage:	4 °C,-20 °C
Storage Comment:	Shipped at 4°C. Store at -20°C for one year. Avoid repeated freeze/thaw cycles.
Expiry Date:	12 months