

Datasheet for ABIN1714774
anti-FTSJ1 antibody (AA 1-110)[Go to Product page](#)

1 Image

Overview

Quantity:	100 µL
Target:	FTSJ1
Binding Specificity:	AA 1-110
Reactivity:	Mouse, Rat
Host:	Rabbit
Clonality:	Polyclonal
Conjugate:	This FTSJ1 antibody is un-conjugated
Application:	ELISA, Immunofluorescence (Paraffin-embedded Sections) (IF (p)), Immunofluorescence (Cultured Cells) (IF (cc)), Immunohistochemistry (Frozen Sections) (IHC (fro)), Immunohistochemistry (Paraffin-embedded Sections) (IHC (p)), Immunocytochemistry (ICC)

Product Details

Immunogen:	KLH conjugated synthetic peptide derived from human FTSJ1
Isotype:	IgG
Cross-Reactivity:	Mouse, Rat
Predicted Reactivity:	Human,Dog,Cow,Sheep,Pig,Horse,Rabbit
Purification:	Purified by Protein A.

Target Details

Target:	FTSJ1
---------	-------

Target Details


Alternative Name:	FTSJ1 (FTSJ1 Products)
Background:	<p>Synonyms: CDLIV, FTSJ 1, FtsJ homolog 1 E. coli, FtsJ homolog 1, JM23, Mental retardation X linked 44, Mental retardation X linked 9, MRX44, MRX9, Putative ribosomal RNA methyltransferase 1, RRMJ1, SPB1, TRM7, RRMJ1_HUMAN.</p> <p>Background: FTSJ1 is a 329 amino acid nucleolar protein belonging to the RImE family and methyltransferase superfamily. Expressed in adult thalamus, hippocampus, amygdala, corpus callosum and caudate nucleus, as well as fetal kidney, lung, liver, brain and lung, FTSJ1 plays a role in rRNA modification and processing. FTSJ1 exists as multiple spliced isoforms which are encoded by a gene located on human chromosome Xp11.23. Notably, defects in the gene encoding FTSJ1 are the cause of mental retardation X-linked type 44 (MRX44) and nonsyndromic X-linked mental retardation (MRX9).</p>

Application Details

Application Notes:	ELISA 1:500-1000 IHC-P 1:200-400 IHC-F 1:100-500 IF(IHC-P) 1:50-200 IF(IHC-F) 1:50-200 IF(ICC) 1:50-200 ICC 1:100-500
Restrictions:	For Research Use only

Handling

Format:	Liquid
Concentration:	1 µg/µL
Buffer:	0.01M TBS(pH 7.4) with 1 % BSA, 0.02 % Proclin300 and 50 % Glycerol.
Preservative:	ProClin
Precaution of Use:	This product contains ProClin: a POISONOUS AND HAZARDOUS SUBSTANCE, which should be handled by trained staff only.
Storage:	4 °C,-20 °C
Storage Comment:	Shipped at 4°C. Store at -20°C for one year. Avoid repeated freeze/thaw cycles.
Expiry Date:	12 months



Immunohistochemistry (Paraffin-embedded Sections)

Image 1. Paraformaldehyde-fixed, paraffin embedded rat lung, Antigen retrieval by boiling in sodium citrate buffer (pH6) for 15min, Block endogenous peroxidase by 3% hydrogen peroxide for 30 minutes, Blocking buffer (normal goat serum) at 37°C for 20min, Antibody incubation with FTSJ1 Polyclonal Antibody, Unconjugated at 1:500 overnight at 4°C, followed by a conjugated secondary and DAB staining.