# antibodies -online.com





## anti-HES3 antibody (AA 11-100)



( )	ve	K\ /		A .
	$\cup$	1 V/	-	V۷

Quantity:	100 μL
Target:	HES3
Binding Specificity:	AA 11-100
Reactivity:	Human
Host:	Rabbit
Clonality:	Polyclonal
Conjugate:	This HES3 antibody is un-conjugated
Application:	Immunofluorescence (Cultured Cells) (IF (cc)), Immunofluorescence (Paraffin-embedded Sections) (IF (p)), ELISA, Immunohistochemistry (Frozen Sections) (IHC (fro)), Immunohistochemistry (Paraffin-embedded Sections) (IHC (p)), Immunocytochemistry (ICC)

#### Product Details

Immunogen:	KLH conjugated synthetic peptide derived from human HES3
Isotype:	IgG
Cross-Reactivity:	Human
Predicted Reactivity:	Mouse,Rat,Dog,Cow,Sheep,Pig,Horse,Rabbit
Purification:	Purified by Protein A.

#### Target Details

Target: HES	33					

#### Target Details

Alternative Name:	HES3 (HES3 Products)
Background:	Synonyms: bHLHb43, Class B basic helix loop helix protein 43, Hairy and enhancer of split 3,
	Transcription factor HES 3,
	Background: The Drosophila hairy and Enhancer of split genes encode basic helix-loop-helix
	(bHLH) transcriptional repressors that function in the Notch signaling pathway and control
	segmentation and neural development during embryogenesis. The mammalian homologues of
	Drosophila hairy and Enhancer of split are the HES gene family members, HES1-6, which also
	encode bHLH transcriptional repressors that regulate myogenesis and neurogenesis. The HES
	family members form a complex with TLE, the mammalian homologue of Groucho, and this
	interaction is mediated by the carboxy terminal WRPW motif of the HES proteins. The HES/TLE
	complex functions by directly binding to DNA, instead of interfering with activator proteins.
	Most HES family members, including HES1 and HES5, preferentially bind to the N box
	(CACNAG) as opposed to the E box (CANNTG). HES2 binds to both N and E box sites, while
	HES6 does not bind DNA. Rather, HES6 inhibits HES1 activity, thereby promoting transcription.
	HES1 and HES2 are expressed in a variety of adult and embryonic tissues. HES3 is expressed
	exclusively in cerebellar Purkinje cells, and HES5 is found solely in the nervous system. HES6 is
	produced in brain as well as in the limb buds of developing embryos.
Gene ID:	390992
Application Details	
Application Notes:	ELISA 1:500-1000
	IHC-P 1:200-400
	IHC-F 1:100-500
	IF(IHC-P) 1:50-200
	IF(IHC-F) 1:50-200
	IF(ICC) 1:50-200
	ICC 1:100-500
Restrictions:	For Research Use only
Handling	
9	
Format:	Liquid
-	Liquid 1 μg/μL

### Handling

Preservative:	ProClin
Precaution of Use:	This product contains ProClin: a POISONOUS AND HAZARDOUS SUBSTANCE, which should be handled by trained staff only.
Storage:	4 °C,-20 °C
Storage Comment:	Shipped at 4°C. Store at -20°C for one year. Avoid repeated freeze/thaw cycles.
Expiry Date:	12 months