

Datasheet for ABIN1714926
anti-GABBR2 antibody (pSer884)



[Go to Product page](#)

Overview

Quantity:	100 µL
Target:	GABBR2
Binding Specificity:	pSer884
Reactivity:	Rat, Mouse
Host:	Rabbit
Clonality:	Polyclonal
Conjugate:	This GABBR2 antibody is un-conjugated
Application:	Western Blotting (WB), ELISA, Immunofluorescence (Cultured Cells) (IF (cc)), Immunofluorescence (Paraffin-embedded Sections) (IF (p)), Immunocytochemistry (ICC), Immunohistochemistry (Paraffin-embedded Sections) (IHC (p)), Immunohistochemistry (Frozen Sections) (IHC (fro))

Product Details

Immunogen:	KLH conjugated synthetic phosphopeptide derived from human GABBR2 around the phosphorylation site of Ser884
Isotype:	IgG
Cross-Reactivity:	Mouse, Rat
Predicted Reactivity:	Human,Dog,Cow,Sheep,Pig,Horse,Rabbit,Guinea Pig
Purification:	Purified by Protein A.

Target Details

Target:	GABBR2
Alternative Name:	GABBR2 (GABBR2 Products)
Background:	Synonyms: GABA B Receptor 2 phospho S884, p-GABA B Receptor 2 phospho S884, p-GABBR2 Ser884 , GABBR2, GB2, G protein coupled receptor 51, G-protein coupled receptor 51, GAB B R2, GABA B R2, GABA B receptor 2, GABA B RECEPTOR, GABA-B receptor 2, GABA-B-R2, GABA-BR2, GABA[[B]]R2, GABAB R2, GABABR 2, GABABR2, GABB R2, GABBR 2, Gabbr2, GABR2_HUMAN, Gamma aminobutyric acid GABA B receptor 2, Gamma aminobutyric acid B receptor 2, Gamma aminobutyric acid GABA B receptor 2, Gamma aminobutyric acid type B receptor subunit 2, Gamma-aminobutyric acid type B receptor subunit 2, Gb 2, Gb2, GH07312, GPR 51, GPR51, GPRC 3B, GPRC3B, HG 20, HG20, HRIHFB2099, Metabotropic GABA B receptor subtype 2, OTTHUMP00000021776, OTTHUMP00000063797, R2 SUBUNIT, BcDNA:GH07312, CG6706, CT20836, D GABA[[B]]R2, D Gaba2, FLJ36928. Background: The multi-pass membrane protein encoded by this gene belongs to the G-protein coupled receptor 3 family and GABA-B receptor subfamily. The GABA-B receptors inhibit neuronal activity through G protein-coupled second-messenger systems, which regulate the release of neurotransmitters, and the activity of ion channels and adenylyl cyclase. This receptor subunit forms an active heterodimeric complex with GABA-B receptor subunit 1, neither of which is effective on its own. Allelic variants of this gene have been associated with nicotine dependence.[provided by RefSeq, Jan 2010].
Gene ID:	9568
Pathways:	cAMP Metabolic Process

Application Details

Application Notes:	WB 1:300-5000 ELISA 1:500-1000 IHC-P 1:200-400 IHC-F 1:100-500 IF(IHC-P) 1:50-200 IF(IHC-F) 1:50-200 IF(ICC) 1:50-200 ICC 1:100-500
Restrictions:	For Research Use only

Handling

Format:	Liquid
Concentration:	1 µg/µL
Buffer:	0.01M TBS(pH 7.4) with 1 % BSA, 0.02 % Proclin300 and 50 % Glycerol.
Preservative:	ProClin
Precaution of Use:	This product contains ProClin: a POISONOUS AND HAZARDOUS SUBSTANCE, which should be handled by trained staff only.
Storage:	4 °C,-20 °C
Storage Comment:	Shipped at 4°C. Store at -20°C for one year. Avoid repeated freeze/thaw cycles.
Expiry Date:	12 months