antibodies - online.com







anti-GALE antibody (AA 21-120)



()	11/0	K\ /	iew
	\cup	'I V/I	$I \cap VV$

Quantity:	100 μL
Target:	GALE
Binding Specificity:	AA 21-120
Reactivity:	Mouse
Host:	Rabbit
Clonality:	Polyclonal
Conjugate:	This GALE antibody is un-conjugated
Application:	Western Blotting (WB), ELISA, Immunofluorescence (Cultured Cells) (IF (cc)), Immunofluorescence (Paraffin-embedded Sections) (IF (p)), Immunohistochemistry (Paraffin-embedded Sections) (IHC (p)), Immunohistochemistry (Frozen Sections) (IHC (fro)), Immunocytochemistry (ICC)

Product Details

Immunogen:	KLH conjugated synthetic peptide derived from human GALE/Galactowaldenase
Isotype:	IgG
Cross-Reactivity:	Mouse
Predicted Reactivity:	Human,Rat,Pig,Horse
Purification:	Purified by Protein A.

Target Details

Target: GALE

Target Details

Alternative Name:	GALE/Galactowaldenase (GALE Products)	
Background:	Synonyms: FLJ95174, FLJ97302, Galactose 4 epimerase UDP, Galactowaldenase, galE,	
васкугочни.	GALE_HUMAN, OTTHUMP0000002991, OTTHUMP0000002994, OTTHUMP0000037931,	
	OTTHUMP00000044857, SDR1E1, short chain dehydrogenase/reductase family 1E member 1,	
	UDP galactose 4 epimerase, UDP glucose 4 epimerase, UDP-galactose 4-epimerase, UDP-	
	glucose 4-epimerase.	
	Background: GALE is a 348 amino acid protein that functions as the third enzyme in the Leloir	
	pathway of galactose metabolism. A member of the sugar epimerase family, GALE exists as a	
	homodimer, binds FAD as a cofactor and catalyzes the epimerization of UDP-N-	
	acetylglucosamine to UDP-N-acetylgalactosamine and UDP-glucose to UDP-galactose. The	
	gene encoding GALE maps to human chromosome 1p36.11 and mutations in this gene lead to	
	the development of complex disorder known as epimerase-deficiency galactosemia (EDG) or	
	galactosemia type 3, which is characterized by mental retardation, liver damage, cataracts and	
	deafness.	
Gene ID:	2582	
Pathways:	Response to Water Deprivation, Cellular Glucan Metabolic Process	
Application Details		
Application Notes:	WB 1:300-5000	
	ELISA 1:500-1000	
	IHC-P 1:200-400	
	IHC-F 1:100-500	
	IF(IHC-P) 1:50-200	
	IF(IHC-F) 1:50-200	
	IF(ICC) 1:50-200	
	ICC 1:100-500	
Restrictions:	For Research Use only	
Handling		
Format:	Liquid	
Concentration:	1 μg/μL	
Buffer:	0.01M TBS(pH 7.4) with 1 % BSA, 0.02 % Proclin300 and 50 % Glycerol.	

Handling

Preservative:	ProClin	
Precaution of Use:	This product contains ProClin: a POISONOUS AND HAZARDOUS SUBSTANCE, which should be handled by trained staff only.	
Storage:	4 °C,-20 °C	
Storage Comment:	nent: Shipped at 4°C. Store at -20°C for one year. Avoid repeated freeze/thaw cycles.	
Expiry Date:	12 months	