antibodies -online.com





Datasheet for ABIN1714996

anti-DAZ4 antibody (AA 51-150)



Overview

Quantity:	100 μL
Target:	DAZ4
Binding Specificity:	AA 51-150
Reactivity:	Mouse
Host:	Rabbit
Clonality:	Polyclonal
Conjugate:	This DAZ4 antibody is un-conjugated
Application:	Western Blotting (WB), ELISA, Immunofluorescence (Cultured Cells) (IF (cc)), Immunofluorescence (Paraffin-embedded Sections) (IF (p)), Immunohistochemistry (Paraffin-embedded Sections) (IHC (p)), Immunohistochemistry (Frozen Sections) (IHC (fro)), Immunocytochemistry (ICC)

Product Details

Immunogen:	KLH conjugated synthetic peptide derived from human DAZ4
Isotype:	IgG
Cross-Reactivity:	Mouse
Predicted Reactivity:	Human,Rat,Dog,Cow,Sheep,Pig,Horse
Purification:	Purified by Protein A.

Target Details

Target:	DAZ4			
---------	------	--	--	--

Target Details

Alternative Name:	DAZ4 (DAZ4 Products)
Background:	Synonyms: DAZ4, Deleted in azoospermia 4, Deleted in azoospermia protein 4.
	Background: Spermatogenesis is the process by which male spermatogonia develop into
	mature spermatozoa. DAZ (deleted in azoospermia) are RNA-binding proteins that play an
	essential role in spermatogenesis. DAZ proteins influence the first stages of spermatogenesis
	and the maintenance of germ cell populations. DAZ proteins (DAZ1, DAZ2, DAZ3, DAZ4 and
	DAZ5) are encoded by separate genes on chromosome Y, each of which contain an AZFc
	domain in their coding region. DAZ proteins localize to the nucleus of spermatogonia, but
	relocate to the cytoplasm during meiosis. DAZ proteins contain an RRM (RNA recognition
	motif) domain that may regulate mRNA translation by binding to the 3?UTR. Deletions in the
	genes encoding DAZ proteins may cause azoospermia or oligospermia which can lead to male
	infertility. DAZ4 (deleted in azoospermia 4), also known as pDP1680 or pDP1681, is a 579
	amino acid testis specific protein that contains nine DAZ-like domains and two RNA recognition
	motifs (RRM). DAZ4 exists as two alternatively spliced isoforms.
Application Details	
Application Notes:	WB 1:300-5000
	ELISA 1:500-1000
	IHC-P 1:200-400
	IHC-F 1:100-500
	IF(IHC-P) 1:50-200
	IF(IHC-F) 1:50-200
	IF(ICC) 1:50-200
	ICC 1:100-500
Restrictions:	For Research Use only
Handling	
Format:	Liquid
Concentration:	1 μg/μL
Buffer:	0.01M TBS(pH 7.4) with 1 % BSA, 0.02 % Proclin300 and 50 % Glycerol.
Preservative:	ProClin
Precaution of Use:	This product contains ProClin: a POISONOUS AND HAZARDOUS SUBSTANCE, which should be
	handled by trained staff only.

Handling

Storage:	4 °C,-20 °C	
Storage Comment:	Shipped at 4°C. Store at -20°C for one year. Avoid repeated freeze/thaw cycles.	
Expiry Date:	12 months	