

Datasheet for ABIN1715081
anti-FANCD2 antibody (pSer1404)



[Go to Product page](#)

Overview

Quantity:	100 µL
Target:	FANCD2
Binding Specificity:	pSer1404
Reactivity:	Human
Host:	Rabbit
Clonality:	Polyclonal
Conjugate:	This FANCD2 antibody is un-conjugated
Application:	ELISA, Immunofluorescence (Cultured Cells) (IF (cc)), Immunofluorescence (Paraffin-embedded Sections) (IF (p)), Immunohistochemistry (Paraffin-embedded Sections) (IHC (p)), Flow Cytometry (FACS), Immunocytochemistry (ICC), Immunohistochemistry (Frozen Sections) (IHC (fro))

Product Details

Immunogen:	KLH conjugated synthetic phosphopeptide derived from human FANCD2 around the phosphorylation site of Ser1404
Isotype:	IgG
Cross-Reactivity:	Human
Predicted Reactivity:	Cow,Sheep,Pig,Horse,Rabbit
Purification:	Purified by Protein A.

Target Details

Target:	FANCD2
Alternative Name:	FANCD2 (FANCD2 Products)
Background:	<p>Synonyms: FANCD2 phospho S1404, FANCD2 phospho Ser1404, p-FANCD2 phospho S1404, FA 4, FA D2, FA4, FAC D2, FACD 2, FACD, FACD2, FAD, FAD2, FANC D2, FANCD 2, FANCD, Protein FACD2, Type 4 Fanconi pancytopenia, FACD2_HUMAN, Fanconi anemia group D2 protein, Protein FACD2.</p> <p>Background: Fanconi Anemia (FANC) is a human autosomal-recessive cancer susceptibility disorder characterized by congenital defects, progressive bone marrow failure, and cellular hypersensitivity to mitomycin C (MMC). The FANC subunit D2 protein is vital for cellular resistance to DNA cross-linking and the arrest of DNA synthesis after ionizing radiation. DNA damage activates the monoubiquitination of FANC D2, targeting nuclear foci containing the BRCA 1 protein.</p>
Gene ID:	2177
Pathways:	DNA Damage Repair

Application Details

Application Notes:	ELISA 1:500-1000 FCM 1:20-100 IHC-P 1:200-400 IHC-F 1:100-500 IF(IHC-P) 1:50-200 IF(IHC-F) 1:50-200 IF(ICC) 1:50-200 ICC 1:100-500
Restrictions:	For Research Use only
Handling	
Format:	Liquid
Concentration:	1 µg/µL
Buffer:	0.01M TBS(pH 7.4) with 1 % BSA, 0.02 % Proclin300 and 50 % Glycerol.
Preservative:	ProClin
Precaution of Use:	This product contains ProClin: a POISONOUS AND HAZARDOUS SUBSTANCE, which should be

Handling

handled by trained staff only.

Storage: 4 °C,-20 °C

Storage Comment: Shipped at 4°C. Store at -20°C for one year. Avoid repeated freeze/thaw cycles.

Expiry Date: 12 months