antibodies - online.com







anti-IL-31 antibody (AA 21-120)





Overview

Quantity:	100 μL
Target:	IL-31 (IL31)
Binding Specificity:	AA 21-120
Reactivity:	Mouse
Host:	Rabbit
Clonality:	Polyclonal
Conjugate:	This IL-31 antibody is un-conjugated
Application:	Western Blotting (WB), ELISA, Immunocytochemistry (ICC), Immunohistochemistry (Paraffinembedded Sections) (IHC (p)), Immunofluorescence (Cultured Cells) (IF (cc)), Immunofluorescence (Paraffin-embedded Sections) (IF (p)), Immunohistochemistry (Frozen Sections) (IHC (fro))

Product Details

Immunogen:	KLH conjugated synthetic peptide derived from mouse IL-31
Isotype:	IgG
Cross-Reactivity:	Mouse
Predicted Reactivity:	Rat
Purification:	Purified by Protein A.

Target Details

Target: IL-31 (IL31)


Target Details

9 - 1 - 1 - 1 - 1 - 1	
Alternative Name:	IL-31 (IL31 Products)
Background:	Synonyms: Interleukin-31, 1700013B14RIK, IL31, interleukin-31 precursor.
	Background: Interleukin-31 (IL-31) is a recently discovered T-cell cytokine closely related to IL-6
	type cytokines and is preferentially produced by T helper type 2 cells. IL-31 activity is mediated
	through the ligand-induced oligomerization of a dimeric receptor complex containing IL-31
	receptor A and oncostatin M receptor. In response to IL-31 binding, these proteins activate the
	JAK/STAT and the AKT signaling pathways. RNA levels of IL-31 receptor A and oncostatin M
	receptor are induced in activated monocytes but are expressed constitutively in epithelial cells
	IL-31, when overexpressed in transgenic mice, results in the development of pruritis, alopecia
	and skin lesions, and in humans may result in atopic dermatitis, suggesting that IL-31 may
	represent a novel target for antipruritic drug development.
Gene ID:	76399
Application Details	
Application Notes:	WB 1:300-5000
	ELISA 1:500-1000
	IHC-P 1:200-400
	IHC-F 1:100-500
	IF(IHC-P) 1:50-200
	IF(IHC-F) 1:50-200
	IF(ICC) 1:50-200
	ICC 1:100-500
Restrictions:	For Research Use only
Handling	
Format:	Liquid
Concentration:	1 μg/μL
Buffer:	0.01M TBS(pH 7.4) with 1 % BSA, 0.02 % Proclin300 and 50 % Glycerol.
Preservative:	ProClin
Precaution of Use:	This product contains ProClin: a POISONOUS AND HAZARDOUS SUBSTANCE, which should be handled by trained staff only.
Storage:	4 °C,-20 °C

Handling


Storage Comment:	Shipped at 4°C. Store at -20°C for one year. Avoid repeated freeze/thaw cycles.
Expiry Date:	12 months

Images



ELISA

Image 1. Antigen: 0.2ug/100ul, Primary: Antiserum, 1:500, 1:1000, 1:2000, 1:4000, 1:8000, 1:16000, 1:32000, Secondary: HRP conjugated Goat-Anti-Rabbit IgG at 1: 5000, TMB staining, Read the data in MicroplateReader by 450nm



ELISA

Image 2. Antigen: 2 μ g/100 μ L Primary: Antiserum, 1:500, 1:1000, 1:2000, 1:4000, 1:8000, 1:16000, 1:32000; Secondary: HRP conjugated Rabbit Anti-Goat IgG at 1: 5000; TMB staining Read the data in Microplate Reader by 450nm.