



[Go to Product page](#)

Datasheet for ABIN1724833

anti-FGF4 antibody (AA 62-123)

4 Images

2 Publications

Overview

Quantity:	100 µL
Target:	FGF4
Binding Specificity:	AA 62-123
Reactivity:	Human
Host:	Mouse
Clonality:	Monoclonal
Conjugate:	This FGF4 antibody is un-conjugated
Application:	ELISA

Product Details

Immunogen:	Purified recombinant fragment of human FGF4 (AA: 62-123) expressed in E. coli.
Clone:	2D7D5
Isotype:	IgG1
Purification:	purified

Target Details

Target:	FGF4
Alternative Name:	FGF4 (FGF4 Products)
Background:	Description: The protein encoded by this gene is a member of the fibroblast growth factor (FGF) family. FGF family members possess broad mitogenic and cell survival activities and are

Target Details

involved in a variety of biological processes including embryonic development, cell growth, morphogenesis, tissue repair, tumor growth and invasion. This gene was identified by its oncogenic transforming activity. This gene and FGF3, another oncogenic growth factor, are located closely on chromosome 11. Co-amplification of both genes was found in various kinds of human tumors. Studies on the mouse homolog suggested a function in bone morphogenesis and limb development through the sonic hedgehog (SHH) signaling pathway.
Aliases: HST, KFGF, HST-1, HSTF1, K-FGF, HBGF-4

Molecular Weight: 22 kDa

Gene ID: 2249

HGNC: 2249

Pathways: [RTK Signaling](#), [Fc-epsilon Receptor Signaling Pathway](#), [EGFR Signaling Pathway](#), [Neurotrophin Signaling Pathway](#), [Stem Cell Maintenance](#)

Application Details

Application Notes: ELISA: 1:10000

Restrictions: For Research Use only

Handling

Format: Liquid

Buffer: Ascitic fluid containing 0.03 % sodium azide.

Preservative: Sodium azide

Precaution of Use: This product contains Sodium azide: a POISONOUS AND HAZARDOUS SUBSTANCE which should be handled by trained staff only.

Storage: 4 °C/-20 °C


Storage Comment: 4°C, -20°C for long term storage

Publications

Product cited in: Galati, Magdinier, Colasanti, Bauwens, Pinte, Ricordy, Giraud-Panis, Pusch, Savino, Cacchione, Gilson: "TRF2 controls telomeric nucleosome organization in a cell cycle phase-dependent manner." in: **PLoS ONE**, Vol. 7, Issue 4, pp. e34386, (2012) ([PubMed](#)).


Diehl, Idowu, Kimmelshue, York, Jackson-Cook, Turner, Holt, Elmore: "Elevated TRF2 in advanced breast cancers with short telomeres." in: **Breast cancer research and treatment**, Vol. 127, Issue 3, pp. 623-30, (2011) ([PubMed](#)).

Images




ELISA

Image 1. Black line: Control Antigen (100 ng), Purple line: Antigen(10 ng), Blue line: Antigen (50 ng), Red line: Antigen (100 ng),



Flow Cytometry

Image 2. Flow cytometric analysis of NIH/3T3 cells using FGF4 mouse mAb (green) and negative control (red).



Western Blotting

Image 3. Western blot analysis using FGF4 mAb against human FGF4 recombinant protein. (Expected MW is 32.6 kDa)

Images

Please check the [product details page](#) for more images. Overall 4 images are available for ABIN1724833.