
antibodies .- online.com









Go to Product page

	r\/	

Specificity:

Purification:

Cross-Reactivity:

Quantity:	100 μg	
Target:	VDAC1	
Binding Specificity:	AA 1-283	
Reactivity:	Human	
Host:	Mouse	
Clonality:	Monoclonal	
Conjugate:	This VDAC1 antibody is conjugated to PE	
Application:	Western Blotting (WB), Immunohistochemistry (IHC), Immunocytochemistry (ICC), Immunofluorescence (IF)	
Product Details		
Immunogen:	Fusion protein amino acids 1-283 (full-length) of human VDAC1. Mouse: 98% identity (279/283 amino acids identical). Rat: 98% identity (279/283 amino acids identical) >60% identity with VDAC2 and VDAC3.	
Clone:	S152B-23	
Isotype:	IgG2a	


Human, Mouse, Rat


Protein G Purified


Detects ~30 kDa. Does not cross-react with VDAC2 or VDAC3 (based on KO validation results).

Target Details

Target Details			
Target:	VDAC1		
Alternative Name:	VDAC1 (VDAC1 Products)		
Background:	Voltage-dependent anion-selective channel protein 1 (also known as VDAC, VDAC1 or outer		
	mitochondrial membrane protein porin 1) is the the outer mitochondrial membrane receptor fo		
	hexokinase and BCL2L1. VDAC forms a channel through the mitochondrial membrane and is		
	involved in small molecule diffusion, cell volume regulation and apoptosis. VDAC may		
	participate in the formation of the permeability transition pore complex (PTPC), which is		
	responsible for the release of mitochondrial products that triggers apoptosis.		
Gene ID:	7416		
UniProt:	P21796		
Application Details			
Application Notes:	• WB (1:1000)		
	optimal dilutions for assays should be determined by the user.		
Comment:	1 μg/ml of ABIN1741231 was sufficient for detection of VDAC1 in 20 μg of rat brain lysate by		
	colorimetric immunoblot analysis using Goat anti-mouse IgG:HRP as the secondary antibody.		
Restrictions:	For Research Use only		
Handling			
Format:	Liquid		
Concentration:	1 mg/mL		
Buffer:	PBS pH 7.4, 50 % glycerol, 0.1 % sodium azide, Storage buffer may change when conjugated		
Preservative:	Sodium azide		
Precaution of Use:	This product contains Sodium azide: a POISONOUS AND HAZARDOUS SUBSTANCE which		
	should be handled by trained staff only.		
Storage:	4 °C		
Storage Comment:	Conjugated antibodies should be stored at 4°C		







Immunocytochemistry

Image 1. Immunocytochemistry/Immunofluorescence analysis using Mouse Anti-VDAC1 Monoclonal Antibody, Clone S152B-23 (ABIN1741231). Tissue: Neuroblastoma cells (SH-SY5Y). Species: Human. Fixation: 4 % PFA for 15 min. Primary Antibody: Mouse Anti-VDAC1 Monoclonal Antibody (ABIN1741231) at 1:100 for overnight at 4 °C with slow rocking. Secondary Antibody: AlexaFluor 488 at 1:1000 for 1 hour at RT. Counterstain: Phalloidin-iFluor 647 (red) F-Actin stain, Hoechst (blue) nuclear stain at 1:800, 1.6 mM for 20 min at RT. (A) Hoechst (blue) nuclear stain. (B) Phalloidin-iFluor 647 (red) F-Actin stain. (C) VDAC1 Antibody (D) Composite.

Western Blotting

Image 2. Western Blot analysis of Rat Brain Membrane showing detection of ~30 kDa VDAC1 protein using Mouse Anti-VDAC1 Monoclonal Antibody, Clone S152B-23 . Lane 1: Molecular Weight Ladder. Lane 2: Rat Brain Membrane. Load: 15 μg. Block: 2% BSA and 2% Skim Milk in 1X TBST. Primary Antibody: Mouse Anti-VDAC1 Monoclonal Antibody at 1:200 for 16 hours at 4°C. Secondary Antibody: Goat Anti-Mouse IgG: HRP at 1:1000 for 1 hour RT. Color Development: ECL solution for 6 min in RT. Predicted/Observed Size: ~30 kDa.

Immunofluorescence (fixed cells)

Image 3. Immunocytochemistry/Immunofluorescence analysis using Mouse Anti-VDAC1 Monoclonal Antibody, Clone S152B-23. Tissue: Neuroblastoma cell line (SK-N-BE). Species: Human. Fixation: 4% Formaldehyde for 15 min at RT. Primary Antibody: Mouse Anti-VDAC1 Monoclonal Antibody at 1:100 for 60 min at RT. Secondary Antibody: Goat Anti-Mouse ATTO 488 at 1:100 for 60 min at RT. Counterstain: Phalloidin Texas Red F-Actin stain; DAPI (blue)

nuclear stain at 1:1000; 1:5000 for 60 min RT, 5 min RT.

Localization: Mitochondrion, Mitochondrion Outer

Membrane, Nucleus. Magnification: 60X. (A) DAPI (blue)

nuclear stain (B) Phalloidin Texas Red F-Actin stain (C)

VDAC1 Antibody (D) Composite.

Please check the product details page for more images. Overall 4 images are available for ABIN1741231.