# antibodies -online.com





## anti-APBA1 antibody (AA 2-265)

3 Images



Publication



Go to Product page

_					
U	V	er	VI	е	W

Quantity:	50 μg
Target:	APBA1
Binding Specificity:	AA 2-265
Reactivity:	Mouse, Rat
Host:	Rabbit
Clonality:	Polyclonal
Application:	Western Blotting (WB), Immunohistochemistry (IHC), Immunocytochemistry (ICC)

## **Product Details**


Immunogen:	Recombinant rat Mint 1 (aa 2-265).	
Specificity:	Specific for Mint 1.	
Purification: Affinity purified with the immunogen. Rabbit serum albumin was added for stabilizat		

## **Target Details**

Target:	APBA1	
Alternative Name:	Mint 1 (APBA1 Products)	
Background:	Synonyms: X11-like protein	
Pathways:	Synaptic Vesicle Exocytosis, Dicarboxylic Acid Transport	

## **Application Details**

Application Notes:	WB: 1 : 1000 (AP staining)	
	IP: not tested yet	
	ICC: 1: 200 up to 1: 500	
	IHC: 1: 200	
Restrictions:	For Research Use only	
Handling		
Format:	Lyophilized	
Reconstitution:	For reconstitution add 50 $\mu$ L H2O to get a 1mg/ml solution of antibody in PBS. Then aliquot and store at -20 $^{\circ}$ C until use.	
Buffer:	PBS	
Handling Advice:	Affinity purified antibodies are less robust than antisera, since protease inhibitors are also	
	removed during purification. Hence, storage at 4 °C for prolonged periods (i.e. several weeks), is not recommended.	
Storage:	-20 °C	
Storage Comment:	Unlabeled lyophilized antibodies are stable in this form without loss of quality at ambient	
	temperatures for several weeks or even months. They can be stored at 4°C for several years.	
	Lyophilized antibodies must not be stored in the freezer, they may be destroyed!	
Publications		
Product cited in:	Boyken, Grønborg, Riedel, Urlaub, Jahn, Chua: "Molecular profiling of synaptic vesicle docking	
	sites reveals novel proteins but few differences between glutamatergic and GABAergic	
	synapses." in: <b>Neuron</b> , Vol. 78, Issue 2, pp. 285-97, (2013) (PubMed).	




### **Immunohistochemistry**

**Image 1.** Indirect immunostaining of PFA fixed mouse brain section with anti-mint 1 (dilution 1 : 200; red) and mouse anti-synapsin (cat. no. 106 001; dilution 1 : 500; green). Mint 1-positive structures in the accessory olfactory bulb (AOB) have been revealed with fluorochromated secondary antibodies. Nuclei have been visualized by DAPI staining (blue).



#### **Western Blotting**

**Image 2.** dilution: 1 : 1000, sample: rat hippocampus homogenate



#### **Immunocytochemistry**

**Image 3.** Indirect immunostaining of rat hippocampus neurons with anti-Mint 1 (dilution 1 : 200; red) and anti-synapsin (cat. no. 106 001; dilution 1 : 500, green).