

Datasheet for ABIN1742454
anti-GABRA2 antibody (AA 29-37)[2 Images](#)[2 Publications](#)[Go to Product page](#)

Overview

Quantity:	100 µL
Target:	GABRA2
Binding Specificity:	AA 29-37
Reactivity:	Human, Mouse, Rat
Host:	Guinea Pig
Clonality:	Polyclonal
Application:	Western Blotting (WB), Immunocytochemistry (ICC), Immunoprecipitation (IP)

Product Details

Immunogen:	Synthetic peptide (aa 29-37 of rat GABA-A receptor alpha2 precursor protein) coupled to key-hole limpet hemocyanin via an added C-terminal cysteine residue.
Specificity:	Specific for GABA-A receptor alpha2.
Purification:	antiserum

Target Details

Target:	GABRA2
Alternative Name:	GABA-A Receptor Alpha2 (GABRA2 Products)
Background:	Synonyms: GABA(A) R alpha2

Application Details


Application Notes:	WB: 1 : 1000 (AP staining) ICC: 1 : 200 up to 1 : 500 IHC: not tested yet
Comment:	WB: This protein aggregates after boiling, making it necessary to run SDS-PAGE with non-boiled samples. ICC: This antibody is suitable for the surface staining of living cells. After washing cells with bound antibodies, they can be fixed and visualized with secondary reagents.
Restrictions:	For Research Use only

Handling

Format:	Liquid
Buffer:	PBS
Handling Advice:	Crude antisera are more robust than monoclonals. With anti-microbials added, they may be stored at 4 °C. Serum does not contain active proteases, in fact, serum itself contains a powerful cocktail of protease inhibitors. Frozen storage (-20 °C), however, is preferable.
Storage:	4 °C/-20 °C
Storage Comment:	Unlabeled antibodies are stable in this form without loss of quality at ambient temperatures for several weeks or even months. They can be stored at 4 °C for several years.


Publications

Product cited in:	Atkinson, Floerchinger, Qiao, Casey, Williamson, Moseley, Stoica, Goddard, Ge, Tullius, Tomlinson: "Donor brain death exacerbates complement-dependent ischemia/reperfusion injury in transplanted hearts." in: Circulation , Vol. 127, Issue 12, pp. 1290-9, (2013) (PubMed).
-------------------	---



Western Blotting

Image 1. dilution: 1 : 1000, sample: unboiled synaptic membrane fraction of rat brain (LP1)



Immunocytochemistry

Image 2. Indirect immunostaining of hippocampus neurons (dilution 1 : 500).