antibodies -online.com





anti-LILRB1 antibody

3 Images

3

Publications



Go to Product page

-						
O	V	e	rv	1	е	W

Quantity:	0.1 mg	
Target:	LILRB1	
Reactivity:	Human	
Host:	Mouse	
Clonality:	Monoclonal	
Conjugate:	This LILRB1 antibody is un-conjugated	
Application:	Western Blotting (WB), Flow Cytometry (FACS), Immunoprecipitation (IP)	

Product Details

Immunogen:	Hairy cell leukaemia cells	
Clone:	GHI-75	
Isotype:	IgG2b kappa	
Specificity:	The mouse monoclonal antibody GHI/75 recognizes an extracellular epitope of CD85j / ILT2, an 110-120 kDa membrane glycoprotein expressed strongly on plasma cells, moderately on circulating B cells, and weakly on monocytes. It is also expressed on T cell and NK cell subsets (variable, individual).	
Cross-Reactivity (Details):	Human	
Purification:	Purified by protein-A affinity chromatography.	
Purity:	> 95 % (by SDS-PAGE)	

Target Details

Target:	LILRB1		
Alternative Name:	CD85j / ILT2 (LILRB1 Products)		
Background:	Leukocyte immunoglobulin like receptor B1,CD85j, also known as ILT-2 (Ig-like transcript 2), LIR-1 (leukocyte Ig-like receptor 1), or LILRB1 (leukocyte Ig-like receptor B1), is a member of Ig superfamily transmembrane glycoproteins named CD85. The CD85j protein is expressed on several types of immune cells (plasma cells, B cells, monocytes, T and NK cell subsets) where binds to MHC class I molecules on antigen-presenting cells and transduces a negative signal that inhibits stimulation of an immune response. It is thought to control inflammatory responses and cytotoxicity to help focus the immune response and limit autoreactivity.,ILT2, LIR1, MIR7, PIRB, ILT-2		
Gene ID:	10859		
UniProt:	Q8NHL6		
Pathways:	Cellular Response to Molecule of Bacterial Origin, Regulation of Leukocyte Mediated Immunity, Positive Regulation of Immune Effector Process, Production of Molecular Mediator of Immune Response		
Application Details			
Application Notes:	Flow cytometry: Recommended dilution: 1-4 μg/mL.		
Restrictions:	For Research Use only		
Handling			
Concentration:	1 mg/mL		
Buffer:	Phosphate buffered saline (PBS), pH 7.4, 15 mM sodium azide		
Preservative:	Sodium azide		
Precaution of Use:	This product contains Sodium azide: a POISONOUS AND HAZARDOUS SUBSTANCE which should be handled by trained staff only.		
Handling Advice:	Do not freeze.		
Storage:	4 °C		
Storage Comment:	Store at 2-8°C. Do not freeze.		


Product cited in:

Lo Monaco, Tremante, Cerboni, Melucci, Sibilio, Zingoni, Nicotra, Natali, Giacomini: "Human leukocyte antigen E contributes to protect tumor cells from lysis by natural killer cells." in: **Neoplasia (New York, N.Y.)**, Vol. 13, Issue 9, pp. 822-30, (2011) (PubMed).

Riteau, Menier, Khalil-Daher, Martinozzi, Pla, Dausset, Carosella, Rouas-Freiss: "HLA-G1 co-expression boosts the HLA class I-mediated NK lysis inhibition." in: **International immunology**, Vol. 13, Issue 2, pp. 193-201, (2001) (PubMed).


Banham, Colonna, Cella, Micklem, Pulford, Willis, Mason: "Identification of the CD85 antigen as ILT2, an inhibitory MHC class I receptor of the immunoglobulin superfamily." in: **Journal of leukocyte biology**, Vol. 65, Issue 6, pp. 841-5, (1999) (PubMed).

Images




Flow Cytometry

Image 1. Flow cytometry surface staining pattern of human peripheral blood stained using anti-human CD85j(GHI/75) purified antibody (concentration in sample 1 μ g/mL) GAM APC.



Flow Cytometry

Image 2. Flow cytometry multicolor surface staining of human lymphocytes stained using anti-human CD85j (GHI/75) purified antibody (concentration in sample 1 μ g/mL) GAM APC and anti-human CD19 (LT19) PE antibody (20 μ L reagent / 100 μ L of peripheral whole blood).



Flow Cytometry

Image 3. Separation of human CD85j positive B cells (red-filled) from neutrophil granulocytes (black-dashed) in flow cytometry analysis (surface staining) of human peripheral whole blood stained using anti-human CD85j(GHI/75) purified antibody (concentration in sample 1 μ g/mL) GAM APC.