

Datasheet for ABIN1774734  
**anti-PDGFR $\alpha$  antibody (FITC)**



[Go to Product page](#)

2 Images

## Overview

Quantity:	100 tests
Target:	PDGFR $\alpha$
Reactivity:	Human
Host:	Mouse
Clonality:	Monoclonal
Conjugate:	This PDGFR $\alpha$ antibody is conjugated to FITC
Application:	Flow Cytometry (FACS)

## Product Details

Purpose:	Anti-Hu CD140a FITC
Immunogen:	CD140a-transfected NIH 3T3 cells
Clone:	16A1
Isotype:	IgG1 kappa
Specificity:	The mouse monoclonal antibody 16A1 recognizes an extracellular epitope of CD140a / PDGFR $\alpha$ , the 170 kDa alpha chain of platelet-derived growth factor receptor, which is widely expressed on a variety of mesenchymal-derived cells and plays pro-proliferative or anti-proliferative roles in various tumours.
Cross-Reactivity (Details):	Human
Purification:	Purified antibody is conjugated with fluorescein isothiocyanate (FITC) under optimum conditions and unconjugated antibody and free fluorochrome are removed by size-exclusion chromatography.

## Target Details


Target:	PDGFRA
Alternative Name:	CD140a ( <a href="#">PDGFRA Products</a> )
Background:	Platelet derived growth factor receptor alpha,CD140a / PDGF-RA (platelet-derived growth factor receptor alpha) is a cell surface receptor for members of platelet-derived growth factor family, whose intracellular part contains a tyrosine kinase domain. CD140a forms homodimers, or heterodimerizes with CD140b / PDGF-RB. Whereas CD140b induces in different cell types their proliferation and migration, the role of CD140a is more controversial, with pro-proliferative or anti-proliferative effects. CD140a has early developmental functions, mediates mesodermal cell migration, and later acts in signaling associated in epithelial-mesenchymal interactions.,PDGFRA, PDGFR2
Gene ID:	5156
UniProt:	<a href="#">P16234</a>
Pathways:	<a href="#">RTK Signaling</a> , <a href="#">Fc-epsilon Receptor Signaling Pathway</a> , <a href="#">EGFR Signaling Pathway</a> , <a href="#">Neurotrophin Signaling Pathway</a> , <a href="#">Platelet-derived growth Factor Receptor Signaling</a>

## Application Details

Application Notes:	Flow cytometry: The reagent is designed for analysis of human blood cells using 4 µL reagent / 100 µL of whole blood or 10 <sup>6</sup> cells in a suspension. The content of a vial (0.4 ml) is sufficient for 100 tests.
Restrictions:	For Research Use only

## Handling

Reconstitution:	No reconstitution is necessary.
Buffer:	Stabilizing phosphate buffered saline (PBS), pH 7.4, 15 mM sodium azide
Preservative:	Sodium azide
Precaution of Use:	This product contains Sodium azide: a POISONOUS AND HAZARDOUS SUBSTANCE which should be handled by trained staff only.
Handling Advice:	<b>Do not freeze.</b> Avoid prolonged exposure to light.
Storage:	4 °C
Storage Comment:	Store at 2-8°C. Protect from prolonged exposure to light. Do not freeze.



### Flow Cytometry

**Image 1.** Separation of cells stained using anti-human CD140a (16A1) FITC antibody (4  $\mu$ L reagent per million cells in 100  $\mu$ L of cell suspension, red-filled) from cells stained using mouse IgG1 isotype control (MOPC-21) FITC antibody (concentration in sample 3  $\mu$ g/mL, same as CD140a FITC concentration, black-dashed) in flow cytometry analysis (surface staining) of PDGF-RA transfected cells.

### Flow Cytometry

**Image 2.** Surface staining of CD140a in CD140a-transfected cells using anti-CD140a (16A1) FITC.