antibodies .- online.com







anti-Apolipoprotein D antibody (AA 70-80)

Images



Overview

Target:

Quantity:	100 μg
Target:	Apolipoprotein D (APOD)
Binding Specificity:	AA 70-80
Reactivity:	Human
Host:	Goat
Clonality:	Polyclonal
Conjugate:	This Apolipoprotein D antibody is un-conjugated
Application:	Western Blotting (WB), Immunohistochemistry (IHC), ELISA
Product Details	
Purpose:	APOD (aa70-80)
Sequence:	ENGKIKVLNQ E
Isotype:	IgG
Cross Departivity:	
Cross-Reactivity:	Human
Purification:	Human Purified from goat serum by ammonium sulphate precipitation followed by antigen affinity
	Purified from goat serum by ammonium sulphate precipitation followed by antigen affinity

Apolipoprotein D (APOD)

Target Details


Alternative Name:	APOD (APOD Products)
Background:	APOD, apolipoprotein D, apo-D
Molecular Weight:	Expected molecular weight: 26 kDa
Gene ID:	347
Gene ID: NCBI Accession:	347 NP_001638

Application Details

Application Notes:	Immunohistochemistry: Paraffin embedded Human Adrenal Gland. Recommended
	concentration: 5 μg/mL.
	Western Blot: Approx 26 kDa band observed in Human Liver and Kidney lysates (calculated MW
	of 21.3 kDa according to NP_001638.1). The observed molecular weight corresponds to the
	glycosylated form. Recommended concentration: 0.3-1 µg/mL. Primary incubation
	Peptide ELISA: antibody detection limit dilution 1:64000.
Restrictions:	For Research Use only


Handling

Format:	Liquid
Concentration:	0.5 mg/mL
Buffer:	Supplied at 0.5 mg/mL in Tris saline, 0.02 % sodium azide, pH 7.3 with 0.5 % bovine serum albumin.
Preservative:	Sodium azide
Precaution of Use:	This product contains Sodium azide: a POISONOUS AND HAZARDOUS SUBSTANCE which should be handled by trained staff only.
Handling Advice:	Minimize freezing and thawing.
Storage:	-20 °C
Storage Comment:	Aliquot and store at -20°C, with minimal freeze/thawing. A working aliquot may be refrigerated at 4°C for a few weeks and still remain viable.



Immunohistochemistry

Image 1. (ABIN1781849) ($5 \,\mu g/mL$) staining of paraffin embedded Human Adrenal Gland. Steamed antigen retrieval with citrate buffer pH 6, AP-staining.



Western Blotting

Image 2. ABIN1781849 (0.3μg/ml) staining of Human Kidney (A) and Liver (B) lysates (35μg protein in RIPA buffer). Detected by chemiluminescence.