

Datasheet for ABIN180947

**anti-PRNP antibody**

2 Images

4 Publications

[Go to Product page](#)

## Overview

Quantity:	0.25 mg
Target:	PRNP
Reactivity:	Sheep
Host:	Mouse
Clonality:	Monoclonal
Conjugate:	This PRNP antibody is un-conjugated
Application:	Immunohistochemistry (Paraffin-embedded Sections) (IHC (p)), Enzyme Immunoassay (EIA)

## Product Details

Immunogen:	Synthetic peptide 146-Re154Re171-182 of ovine PrP
Clone:	2G11
Isotype:	IgG2a
Specificity:	This antibody recognises the scrapie-associated protease-resistant isoform (PrP <sup>Sc</sup> ) of ovine prion protein (PrP). Clone 2G11 specifically recognises the Re151-Re159 sequence and does not recognise the non-pathogenic PrP <sup>C</sup> form.
Characteristics:	Synonyms: Major prion protein, PrP27-30, PrP33-35C, ASCR, PRNP, PRIP
Purification:	Affinity chromatography on Protein G

## Target Details

Target:	PRNP
---------	------

## Target Details

Alternative Name:	CD230 / PrP ( <a href="#">PRNP Products</a> )
Background:	<p>The prion protein is encoded by the PrP gene and is constitutively expressed as the normal cellular isoform PrP<sup>Sc</sup> by many cell types. Transmissible spongiform encephalopathies (TSE) are neurodegenerative diseases in which the accumulation of the abnormal PrP<sup>Sc</sup> isoform in tissues, of the lymphoid and central nervous system, is thought to be a key event. Studies suggest that PrP<sup>Sc</sup> is the causative agent of transmissible spongiform encephalitis (TSE) but this has not been confirmed. Synonyms: ASCR, Major prion protein, PRIP, PRNP, PrP27-30, PrP33-35C</p>
Gene ID:	493887
UniProt:	<a href="#">P23907</a>
Pathways:	<a href="#">Transition Metal Ion Homeostasis</a> , <a href="#">Activated T Cell Proliferation</a>

## Application Details

Application Notes:	<p>ELISA: 1/50 - 1/1000, Tested on peptide. Immunohistochemistry on paraffin sections: suitable for use on tissue fixed in 4 % neutralbuffered formalin. Treatment of tissue sections in 98 % formic acid, for 30 minutes, is recommended prior to pre-treatment with trypsin at 37C for 5 minutes followed by heatmediated retrieval with 10 mM citrate buffer pH 6.1.</p> <p>Other applications not tested.</p> <p>Optimal dilutions are dependent on conditions and should be determined by the user.</p>
Restrictions:	For Research Use only

## Handling

Concentration:	1.0 mg/mL
Buffer:	PBS, pH 7.4 containing 0.09 % Sodium Azide
Preservative:	Sodium azide
Precaution of Use:	This product contains sodium azide: a POISONOUS AND HAZARDOUS SUBSTANCE which should be handled by trained staff only.
Storage:	4 °C/-20 °C
Storage Comment:	<p>Store the antibody undiluted at 2-8 °C for one month or (in aliquots) at -20 °C for longer. Avoid repeated freezing and thawing.</p> <p>Shelf life: one year from despatch.</p>


## Handling

Expiry Date: 12 months

## Publications


Product cited in: Anish, Hossain, Jacobson, Takada: "Characterization of transcription from TATA-less promoters: identification of a new core promoter element XCPE2 and analysis of factor requirements." in: **PLoS ONE**, Vol. 4, Issue 4, pp. e5103, (2009) ([PubMed](#)).

## Images



### Immunohistochemistry (Paraffin-embedded Sections)

**Image 1.** Immunohistochemistry: 1:50-1:300 working dilutions of anti-Prion Protein, clone 2G11 localizes prion protein in rat brain paraffin embedded tissue (High & Low Magnification).



### Western Blotting

**Image 2.** Rat brain lysate was resolved by electrophoresis, transferred to PVDF and probed with anti-Prion Protein, clone 2G11 (1 µg/mL). Proteins were visualized using a goat anti-mouse secondary antibody conjugated to HRP and a chemiluminescence detection system. Arrow indicates Prion Protein (~27 kDa)