.-online.com antibodies

Datasheet for ABIN184778 anti-MAGOH antibody (C-Term)



Overview

Quantity:	100 µg
Target:	MAGOH
Binding Specificity:	C-Term
Reactivity:	Human
Host:	Goat
Clonality:	Polyclonal
Conjugate:	This MAGOH antibody is un-conjugated
Application:	Western Blotting (WB), ELISA

Product Details

Purpose:	MAGOH
Immunogen:	Peptide with sequence C-SLIGLHFKIKPI, from the C Terminus of the protein sequence according to NP_002361.1.
Sequence:	SLIGLHFKIK PI
lsotype:	lgG
Cross-Reactivity:	Cow, Dog, Human, Mouse, Rat
Purification:	Purified from goat serum by ammonium sulphate precipitation followed by antigen affinity chromatography using the immunizing peptide.
Grade:	Verified

Order at www.antibodies-online.com | www.antikoerper-online.de | www.anticorps-enligne.fr | www.antibodies-online.cn International: +49 (0)241 95 163 153 | USA & Canada: +1 877 302 8632 | support@antibodies-online.com Page 1/2 | Product datasheet for ABIN184778 | 12/22/2023 | Copyright antibodies-online. All rights reserved.

Target Details	
Target:	MAGOH
Alternative Name:	MAGOH (MAGOH Products)
Background:	MAGOH, mago-nashi homolog, proliferation-associated (Drosophila), RP5-1024G6.4, MAGOHA
Gene ID:	4116, 17149
NCBI Accession:	NP_002361
Application Details	
Application Notes:	Western Blot: Approx 17 kDa band observed in nuclear lysates of cell line HeLa (calculated MW of 17.2 kDa according to NP_002361.1). Recommended concentration: 1-3 µg/mL. (Data obtained from a previous batch). Peptide ELISA: antibody detection limit dilution 1:2000.
Restrictions:	For Research Use only
Handling	
Format:	Liquid
Concentration:	0.5 mg/mL
Buffer:	Supplied at 0.5 mg/mL in Tris saline, 0.02 % sodium azide, pH 7.3 with 0.5 % bovine serum albumin.
Preservative:	Sodium azide
Precaution of Use:	This product contains Sodium azide: a POISONOUS AND HAZARDOUS SUBSTANCE which should be handled by trained staff only.
Handling Advice:	Minimize freezing and thawing.
Storage:	-20 °C
Storage Comment:	Aliquot and store at -20°C, with minimal freeze/thawing. A working aliquot may be refrigerated at 4°C for a few weeks and still remain viable.