antibodies -online.com





anti-UBE2L3 antibody (C-Term)



Images



Go to Product page

Overview


Quantity:	100 μg
Target:	UBE2L3
Binding Specificity:	C-Term
Reactivity:	Human, Mouse, Rat, Pig
Host:	Goat
Clonality:	Polyclonal
Conjugate:	This UBE2L3 antibody is un-conjugated
Application:	Western Blotting (WB), ELISA, Immunofluorescence (IF)


Product Details

1 Toddet Details	
Purpose:	UBE2L3
Immunogen:	Peptide with sequence C-EFTKKYGEKRPVD, from the C Terminus of the protein sequence according to NP_003338.1.
Sequence:	EFTKKYGEKR PVD
Isotype:	IgG
Cross-Reactivity:	Cow, Human, Mouse, Pig, Rat
Purification:	Purified from goat serum by ammonium sulphate precipitation followed by antigen affinity chromatography using the immunizing peptide.
Grade:	Verified

Target Details


rarget Details	
Target:	UBE2L3
Alternative Name:	UBE2L3 (UBE2L3 Products)
Background:	UBE2L3, ubiquitin-conjugating enzyme E2L 3, UBCH7, E2-F1, L-UBC, UbcM4, ubiquitin carrier
	protein, ubiquitin-conjugating enzyme UBCH7, ubiquitin-protein ligase
Gene ID:	7332, 22195
NCBI Accession:	NP_003338
Application Details	
Application Notes:	Western Blot: Approx 18 kDa band observed in lysates of cell lines HeLa HEK293, HepG2 and
	Jurkat and also in lysates of Human, Mouse and Rat Heart, Skeletal Muscle and Testis and o
	Pig Heart (calculated MW of 17.9 kDa according to Human NP_003338.1 and Mo
	Peptide ELISA: antibody detection limit dilution 1:128000.
Comment:	Immunofluorescence: Strong expression of the protein seen in the cytoplasm. Dividing cells
	appear to show enhanced expression. Recommended concentration: 5µg/ml.
Restrictions:	For Research Use only
Handling	
Format:	Liquid
Concentration:	0.5 mg/mL
Buffer:	Supplied at 0.5 mg/mL in Tris saline, 0.02 % sodium azide, pH 7.3 with 0.5 % bovine serum
	albumin.
Preservative:	Sodium azide
Precaution of Use:	This product contains Sodium azide: a POISONOUS AND HAZARDOUS SUBSTANCE which
	should be handled by trained staff only.
Handling Advice:	Minimize freezing and thawing.
Storage:	-20 °C
Storage Comment:	Aliquot and store at -20°C, with minimal freeze/thawing. A working aliquot may be refrigerate
	at 4°C for a few weeks and still remain viable.






Immunofluorescence

Image 1. Immunofluorescence staining of MCF7 (A) and HeLa (B) cells with 5ug/ml ABIN184900 antibody. Detected with Rabbit anti-goat IgG-Alexa Fluor488 antibody at 1:1000. Nuclei Counterstained with DAPI.



Western Blotting

Image 2. ABIN184900 staining (0.5 ug/ml) of Hela lysate (RIPA buffer, 35 ug total protein per lane). Primary incubated for 1 hour. Detected by western blot using chemiluminescence.



Western Blotting

Image 3. ABIN184900 (0.01μg/ml) staining of HEK293 (A), HepG2 (B) and Jurkat (C) lysates (35μg protein in RIPA buffer). Primary incubation was 1 hour. Detected by chemiluminescence.

Please check the product details page for more images. Overall 5 images are available for ABIN184900.