



[Go to Product page](#)

Datasheet for ABIN184959
anti-TRIM40 antibody (C-Term)

1 Image

Overview

Quantity:	100 µg
Target:	TRIM40
Binding Specificity:	C-Term
Reactivity:	Human
Host:	Goat
Clonality:	Polyclonal
Conjugate:	This TRIM40 antibody is un-conjugated
Application:	Western Blotting (WB), ELISA

Product Details

Purpose:	RNF35 / TRIM40
Immunogen:	C-GVSELLLQPPQKL
Sequence:	GVSELLLQPP QKL
Isotype:	IgG
Cross-Reactivity:	Human
Purification:	Purified from goat serum by ammonium sulphate precipitation followed by antigen affinity chromatography using the immunizing peptide.
Grade:	Verified

Target Details


Target:	TRIM40
Alternative Name:	TRIM40 (TRIM40 Products)
Background:	TRIM40, tripartite motif-containing 40, RNF35, ring finger RNF35, DAAP-200B17.3, ring finger protein 40
Gene ID:	135644
NCBI Accession:	NP_619645

Application Details

Application Notes:	Western Blot: Approx 26 kDa band observed in Human Placenta lysates (calculated MW of 26.1 kDa according to NP_619645.1). Recommended concentration: 0.5-2 µg/mL. Peptide ELISA: antibody detection limit dilution 1:32000.
Restrictions:	For Research Use only

Handling

Format:	Liquid
Concentration:	0.5 mg/mL
Buffer:	Supplied at 0.5 mg/mL in Tris saline, 0.02 % sodium azide, pH 7.3 with 0.5 % bovine serum albumin.
Preservative:	Sodium azide
Precaution of Use:	This product contains Sodium azide: a POISONOUS AND HAZARDOUS SUBSTANCE which should be handled by trained staff only.
Handling Advice:	Minimize freezing and thawing.
Storage:	-20 °C
Storage Comment:	Aliquot and store at -20°C, with minimal freeze/thawing. A working aliquot may be refrigerated at 4°C for a few weeks and still remain viable.



Western Blotting

Image 1. ABIN184959 (0.5µg/ml) staining of Human Placenta lysate (35µg protein in RIPA buffer). Primary incubation was 1 hour. Detected by chemiluminescence.