# antibodies - online.com







# anti-FBXL12 antibody (C-Term)



Image



#### Overview

Quantity:	100 μg
Target:	FBXL12
Binding Specificity:	C-Term
Reactivity:	Human
Host:	Goat
Clonality:	Polyclonal
Conjugate:	This FBXL12 antibody is un-conjugated
Application:	ELISA, Immunohistochemistry (IHC)

#### **Product Details**

Purpose:	FBXL12
Immunogen:	Peptide with sequence RACPKESMDWWM, from the C Terminus of the protein sequence according to NP_060173.1.
Sequence:	RACPKESMDW WM
Isotype:	IgG
Cross-Reactivity:	Dog, Human, Mouse, Rat
Purification:	Purified from goat serum by ammonium sulphate precipitation followed by antigen affinity chromatography using the immunizing peptide.
Grade:	Verified

## **Target Details**

rarget Details	
Target:	FBXL12
Alternative Name:	FBXL12 (FBXL12 Products)
Background:	FBXL12, likely ortholog of mouse f-box and leucine-rich repeat protein 12, FLJ20188,
	hypothetical protein FLJ20188, F-box and leucine-rich repeat protein 12, Fbl12
Gene ID:	54850, 30843
NCBI Accession:	NP_060173
Pathways:	SARS-CoV-2 Protein Interactome
Application Details	
Application Notes:	Immunohistochemistry: In paraffin embedded Human Lung shows staining of epithelial cells in
	the bronchioles and alveoli. Recommended concentration: 4-6 µg/mL.
	Peptide ELISA: antibody detection limit dilution 1:128000.
Restrictions:	For Research Use only
Handling	
Format:	Liquid
Concentration:	0.5 mg/mL
Buffer:	Supplied at 0.5 mg/mL in Tris saline, 0.02 % sodium azide, pH 7.3 with 0.5 % bovine serum
	albumin.
Preservative:	Sodium azide
Precaution of Use:	This product contains Sodium azide: a POISONOUS AND HAZARDOUS SUBSTANCE which
	should be handled by trained staff only.
Handling Advice:	Minimize freezing and thawing.


at 4°C for a few weeks and still remain viable.

Aliquot and store at -20°C, with minimal freeze/thawing. A working aliquot may be refrigerated

-20 °C

Storage:

Storage Comment:



### **Immunohistochemistry**

**Image 1.** ABIN185059 ( $4\mu g/ml$ ) staining of paraffin embedded Human Lung. Steamed antigen retrieval with citrate buffer pH 6, HRP-staining.