

Datasheet for ABIN185060
anti-FAF2 antibody (C-Term)

3 Images

[Go to Product page](#)

Overview

Quantity:	100 µg
Target:	FAF2
Binding Specificity:	C-Term
Reactivity:	Human, Mouse
Host:	Goat
Clonality:	Polyclonal
Conjugate:	This FAF2 antibody is un-conjugated
Application:	Western Blotting (WB), ELISA

Product Details

Purpose:	ETEA / UBXD8
Immunogen:	Peptide with sequence C-HTEVLVQDLTDE, from the C Terminus of the protein sequence according to NP_055428.1.
Sequence:	HTEVLVQDL TDE
Isotype:	IgG
Cross-Reactivity:	Cow, Dog, Human, Mouse, Rat
Purification:	Purified from goat serum by ammonium sulphate precipitation followed by antigen affinity chromatography using the immunizing peptide.
Grade:	Verified

Target Details

Target:	FAF2
Alternative Name:	FAF2 (FAF2 Products)
Background:	FAF2, ETEA, UBXD8, KIAA0887, expressed in T-cells and eosinophils in atopic dermatitis, UBX domain containing 8, Fas associated factor family member 2, UBXN3B, UBX domain protein 3B
Gene ID:	23197, 76577
NCBI Accession:	NP_055428

Application Details

Application Notes:	Western Blot: Approx 55 kDa band observed in lysates of Human and Mouse Thymus and of cell line Jurkat (calculated MW of 52.6 kDa according to NP_055428.1). Recommended concentration: 0.3-1 µg/mL. Peptide ELISA: antibody detection limit dilution 1:16000.
Restrictions:	For Research Use only

Handling

Format:	Liquid
Concentration:	0.5 mg/mL
Buffer:	Supplied at 0.5 mg/mL in Tris saline, 0.02 % sodium azide, pH 7.3 with 0.5 % bovine serum albumin.
Preservative:	Sodium azide
Precaution of Use:	This product contains Sodium azide: a POISONOUS AND HAZARDOUS SUBSTANCE which should be handled by trained staff only.
Handling Advice:	Minimize freezing and thawing.
Storage:	-20 °C
Storage Comment:	Aliquot and store at -20°C, with minimal freeze/thawing. A working aliquot may be refrigerated at 4°C for a few weeks and still remain viable.



Western Blotting

Image 1. ABIN185060 staining (0.3µg/ml) of Jurkat lysate (RIPA buffer, 35µg total protein per lane). Primary incubated for 1 hour. Detected by chemiluminescence.

Western Blotting

Image 2. ABIN185060 staining (1µg/ml) of Human Testis lysate (RIPA buffer, 35µg total protein per lane). Primary incubated for 1 hour. Detected by western blot using chemiluminescence.

Western Blotting

Image 3. ABIN185060 (1µg/ml) staining of Human (A) and Mouse (B) Thymus lysate (35µg protein in RIPA buffer). Primary incubation was 1 hour. Detected by chemiluminescence.