antibodies - online.com







anti-ZBTB32 antibody (C-Term)



Image



Overview

| Quantity: | 100 μg |
|----------------------|---------------------------------------|
| Target: | ZBTB32 |
| Binding Specificity: | C-Term |
| Reactivity: | Human |
| Host: | Goat |
| Clonality: | Polyclonal |
| Conjugate: | This ZBTB32 antibody is un-conjugated |
| Application: | Western Blotting (WB), ELISA |

Product Details

| Purpose: | FAZF / TZFP / ZBTB32 |
|-------------------|---|
| Immunogen: | Peptide with sequence RPSTSPCCPSSSTT, from the C Terminus of the protein sequence according to NP_055198.1. |
| Sequence: | RPSTSPCCPS SSTT |
| Isotype: | IgG |
| Cross-Reactivity: | Human |
| Purification: | Purified from goat serum by ammonium sulphate precipitation followed by antigen affinity chromatography using the immunizing peptide. |
| Grade: | Verified |

Target Details

| rarget Details | |
|---------------------|---|
| Target: | ZBTB32 |
| Alternative Name: | ZBTB32 (ZBTB32 Products) |
| Background: | ZBTB32, TZFP, FAZF, testis zinc finger protein, FAXF, zinc finger and BTB domain containing 32, Rog, ZNF538, OTTHUMP00000045790, repressor of GATA |
| Gene ID: | 27033 |
| NCBI Accession: | NP_055198 |
| Application Details | |
| Application Notes: | Western Blot: A faint approx 55 kDa band observed in Human Bone Marrow lysates (calculated MW of 53.0 kDa according to NP_055198). Recommended concentration: 2-5 µg/mL. Peptide ELISA: antibody detection limit dilution 1:8000. |
| Restrictions: | For Research Use only |
| Handling | |
| Format: | Liquid |
| Concentration: | 0.5 mg/mL |
| Buffer: | Supplied at 0.5 mg/mL in Tris saline, 0.02 % sodium azide, pH 7.3 with 0.5 % bovine serum albumin. |
| Preservative: | Sodium azide |
| Precaution of Use: | This product contains Sodium azide: a POISONOUS AND HAZARDOUS SUBSTANCE which should be handled by trained staff only. |
| Handling Advice: | Minimize freezing and thawing. |
| Storage: | -20 °C |
| Storage Comment: | Aliquot and store at -20°C, with minimal freeze/thawing. A working aliquot may be refrigerated at 4°C for a few weeks and still remain viable. |

250kDa 150kDa 100kDa 75kDa 50kDa 37kDa 25kDa 20kDa 15kDa

Image 1. ABIN185091 (2μg/ml) staining of Human Bone Marrow lysate (35μg protein in RIPA buffer). Primary incubation was 1 hour. Detected by chemiluminescence.