



[Go to Product page](#)

Datasheet for ABIN185694

anti-SNX26 antibody (N-Term)

1 Image

Overview

Quantity:	100 µg
Target:	SNX26
Binding Specificity:	N-Term
Reactivity:	Human
Host:	Goat
Clonality:	Polyclonal
Conjugate:	This SNX26 antibody is un-conjugated
Application:	ELISA, Immunohistochemistry (IHC)

Product Details

Purpose:	SNX26 / TCGAP (N Terminus)
Immunogen:	Peptide with sequence ARSTDSLDPGE-C, from the N Terminus of the protein sequence according to NP_443180.2.
Sequence:	ARSTDSLDPG GE
Isotype:	IgG
Cross-Reactivity:	Human
Purification:	Purified from goat serum by ammonium sulphate precipitation followed by antigen affinity chromatography using the immunizing peptide.
Grade:	Verified

Target Details


Target:	SNX26
Alternative Name:	SNX26 (SNX26 Products)
Background:	SNX26, TCGAP, sorting nexin 26, FLJ39019
Gene ID:	115703
NCBI Accession:	NP_443180

Application Details

Application Notes:	<p>Immunohistochemistry: In paraffin embedded Human Prostate shows cytoplasm staining of secretory cells in the glands. Recommended concentration, 1.25 µg/mL.</p> <p>Western Blot: Preliminary experiments gave an approx 35 kDa band in human brain (hippocampus) lysates after 1 µg/mL antibody staining. Please note that currently we cannot find an explanation in the literature for the band we observe given the calculated s</p> <p>Peptide ELISA: antibody detection limit dilution 1:32000.</p>
Restrictions:	For Research Use only

Handling

Format:	Liquid
Concentration:	0.5 mg/mL
Buffer:	Supplied at 0.5 mg/mL in Tris saline, 0.02 % sodium azide, pH 7.3 with 0.5 % bovine serum albumin.
Preservative:	Sodium azide
Precaution of Use:	This product contains Sodium azide: a POISONOUS AND HAZARDOUS SUBSTANCE which should be handled by trained staff only.
Handling Advice:	Minimize freezing and thawing.
Storage:	-20 °C
Storage Comment:	Aliquot and store at -20°C, with minimal freeze/thawing. A working aliquot may be refrigerated at 4°C for a few weeks and still remain viable.



Immunohistochemistry

Image 1. ABIN185694 (1.25µg/ml) staining of paraffin embedded human prostate. Steamed antigen retrieval with citrate buffer pH 6, AP-staining.