

Datasheet for ABIN1857800
anti-Testosterone antibody (Biotin)



[Go to Product page](#)

Overview

Quantity:	200 µL
Target:	Testosterone
Reactivity:	Various Species
Host:	Rabbit
Clonality:	Polyclonal
Conjugate:	This Testosterone antibody is conjugated to Biotin
Application:	Immunohistochemistry (IHC), Western Blotting (WB), Immunocytochemistry (ICC)

Product Details

Purpose:	Biotin-Linked Polyclonal Antibody to Testosterone (Testo)
Immunogen:	PAA458Ge01 Polyclonal Antibody to Testosterone (Testo)
Isotype:	IgG
Specificity:	The antibody is a rabbit polyclonal antibody raised against Testo. It has been selected for its ability to recognize Testo in immunohistochemical staining and western blotting.
Characteristics:	Chemical Formula: C ₁₉ H ₂₈ O ₂ Molecular Mass: 288.42 g/mol
Purification:	Antigen-specific affinity chromatography followed by Protein A affinity chromatography

Target Details

Target:	Testosterone
---------	--------------

Target Details

Abstract: [Testosterone Products](#)

Target Type: Hormone

Background: T, Testesteron, Testeronun

Application Details

Application Notes: Immunohistochemistry: 5-20 µg/mL Immunocytochemistry: 5-20 µg/mL Optimal working dilutions must be determined by end user.

Comment: The thermal stability is described by the loss rate. The loss rate was determined by accelerated thermal degradation test, that is, incubate the protein at 37°C for 48h, and no obvious degradation and precipitation were observed. The loss rate is less than 5% within the expiration date under appropriate storage condition.

Restrictions: For Research Use only

Handling

Format: Liquid

Concentration: 500 µg/mL

Buffer: 0.01M PBS, pH 7.4, containing 0.05 % Proclin-300, 50 % glycerol.

Preservative: ProClin

Precaution of Use: WARNING: Reagents contain sodium azide. Sodium azide is very toxic if ingested or inhaled. Avoid contact with skin, eyes, or clothing. Wear eye or face protection when handling. If skin or eye contact occurs, wash with copious amounts of water. If ingested or inhaled, contact a physician immediately. Sodium azide yields toxic hydrazoic acid under acidic conditions. Dilute azide-containing compounds in running water before discarding to avoid accumulation of potentially explosive deposits in lead or copper plumbing.

Handling Advice: Avoid repeated freeze/thaw cycles

Storage: 4 °C, -20 °C

Storage Comment: Store at 4°C for frequent use. Stored at -20°C in a manual defrost freezer for two year without detectable loss of activity. Avoid repeated freeze-thaw cycles.

Expiry Date: 12 months