
# antibodies - online.com







# anti-ACVR2A antibody (AA 326-485)

**Images** 



$\sim$			
	IV/E	۱//۱۲	$I \cap V$

Quantity:	100 μL
Target:	ACVR2A
Binding Specificity:	AA 326-485
Reactivity:	Mouse
Host:	Rabbit
Clonality:	Polyclonal
Conjugate:	This ACVR2A antibody is un-conjugated
Application:	Western Blotting (WB), Immunohistochemistry (IHC), Immunocytochemistry (ICC)

#### **Product Details**


Immunogen:	ACVR2A (Lys326-Leu485)	
Isotype:	IgG	
Specificity:	The antibody is a rabbit polyclonal antibody raised against ACVR2A. It has been selected for its ability to recognize ACVR2A in immunohistochemical staining and western blotting.	
Purification:	Antigen-specific affinity chromatography	

#### **Target Details**

Target:	ACVR2A	
Alternative Name:	Activin A Receptor Type II A (ACVR2A) (ACVR2A Products)	
Background:	Alternative Names: ACVR2-A, ACTRII, ACTR-II, ACVR2	


## **Application Details**

1-1		
Application Notes:	<ul> <li>Western blotting: 1:50-400 Immunocytochemistry in formalin fixed cells: 1:50-500         Immunohistochemistry in formalin fixed frozen section: 1:50-500 Immunohistochemistry in paraffin section: 1:10-100 Enzyme-linked Immunosorbent Assay: 1:100-1:5000 Optimal working dilutions must be determined by end user.     </li> </ul>	
Comment:	The thermal stability is described by the loss rate. The loss rate was determined by accelerated	
	thermal degradation test, that is, incubate the protein at 37&degC for 48h, and no obvious degradation and precipitation were observed. The loss rate is less than 5% within the expiration	
	date under appropriate storage condition.	
Restrictions:	For Research Use only	
Handling		
Format:	Liquid	
Concentration:	Lot specific	
Buffer:	PBS, pH 7.4, containing 0.02 % Sodium azide, 50 % glycerol.	
Preservative:	Sodium azide	
Precaution of Use:	WARNING: Reagents contain sodium azide. Sodium azide is very toxic if ingested or inhaled. Avoid contact with skin, eyes, or clothing. Wear eye or face protection when handling. If skin or eye contact occurs, wash with copious amounts of water. If ingested or inhaled, contact a physician immediately. Sodium azide yields toxic hydrazoic acid under acidic conditions. Dilute azide-containing compounds in running water before discarding to avoid accumulation of potentially explosive deposits in lead or copper plumbing.	
Handling Advice:	Avoid repeated freeze-thaw cycles.	
Storage:	4 °C	
Storage Comment:	Store at 2-8 °C for one month. Aliquot and store at -80 °C for 12 months.	
Expiry Date:	12 months	



### **Western Blotting**

Image 1.



#### **Immunohistochemistry**

**Image 2.** Figure.DAB staining on IHC-P. Samples: Mouse Tissue