# antibodies - online.com







# anti-ADAM10 antibody (AA 218-445)



**Images** 



| ( ) | 11/0               | r\ /1      | $\triangle 1 $ |
|-----|--------------------|------------|----------------|
|     | $\lor \lor \vdash$ | $I \vee I$ | ew             |
|     |                    |            |                |

| Quantity:            | 100 μL   |  |
|----------------------|--|--|
| Target:              | ADAM10   |  |
| Binding Specificity: | AA 218-445   |  |
| Reactivity:          | Rat  |  |
| Host:                | Rabbit   |  |
| Clonality:           | Polyclonal   |  |
| Conjugate:           | This ADAM10 antibody is un-conjugated  |  |
| Application:         | Western Blotting (WB), Immunohistochemistry (IHC), Immunocytochemistry (ICC) |  |

#### **Product Details**


| Immunogen:    | ADAM10 (Thr218-Asp445)   |  |
|---------------|--|--|
| Isotype:      | IgG  |  |
| Specificity:  | The antibody is a rabbit polyclonal antibody raised against ADAM10. It has been selected for its ability to recognize ADAM10 in immunohistochemical staining and western blotting. |  |
| Purification: | Antigen-specific affinity chromatography   |  |

#### **Target Details**

| Target:           | ADAM10  |
|-------------------|---|
| Alternative Name: | A Disintegrin And Metalloprotease 10 (ADAM10) (ADAM10 Products)                   |
| Background:       | Alternative Names: CD156-C, CD156c, MADM, Kuz, CDw156, Kuzbanian protein homolog, |


### **Target Details**

| Mammalian disintegrin-metalloprotease  |  |
|--|--|
| Notch Signaling, EGFR Signaling Pathway  |  |
|  |  |
| Western blotting: 1:50-400 Immunocytochemistry in formalin fixed cells: 1:50-500 Immunohistochemistry in formalin fixed frozen section: 1:50-500 Immunohistochemistry in paraffin section: 1:10-100 Enzyme-linked Immunosorbent Assay: 1:100-1:5000 Optimal working dilutions must be determined by end user.  |  |
| The thermal stability is described by the loss rate. The loss rate was determined by accelerated thermal degradation test, that is, incubate the protein at 37&degC for 48h, and no obvious degradation and precipitation were observed. The loss rate is less than 5% within the expiration date under appropriate storage condition.   |  |
| For Research Use only  |  |
|  |  |
| Liquid   |  |
| Lot specific   |  |
| PBS, pH 7.4, containing 0.02 % Sodium azide, 50 % glycerol.  |  |
| Sodium azide   |  |
| WARNING: Reagents contain sodium azide. Sodium azide is very toxic if ingested or inhaled. Avoid contact with skin, eyes, or clothing. Wear eye or face protection when handling. If skin or eye contact occurs, wash with copious amounts of water. If ingested or inhaled, contact a physician immediately. Sodium azide yields toxic hydrazoic acid under acidic conditions. Dilute azide-containing compounds in running water before discarding to avoid accumulation of potentially explosive deposits in lead or copper plumbing. |  |
| Avoid repeated freeze-thaw cycles.   |  |
| 4 °C   |  |
| Store at 2-8 °C for one month. Aliquot and store at -80 °C for 12 months.  |  |
| 12 months  |  |
|  |  |




## Western Blotting

Image 1.



#### **Immunohistochemistry**

**Image 2.** Used in DAB staining on fromalin fixed paraffinembedded kidney tissue



#### **Immunohistochemistry**

Image 3. DAB staining on IHC-P; Samples: Rat Spinal cord Tissue