antibodies -online.com





anti-ADAMTS1 antibody (AA 854-967)

3 Images



Go to Product page

Overview

| Quantity: | 100 μL | |
|----------------------|--|--|
| Target: | ADAMTS1 | |
| Binding Specificity: | AA 854-967 | |
| Reactivity: | Rat | |
| Host: | Rabbit | |
| Clonality: | Polyclonal | |
| Conjugate: | This ADAMTS1 antibody is un-conjugated | |
| Application: | Western Blotting (WB), Immunohistochemistry (IHC), Immunocytochemistry (ICC) | |
| Product Potails | | |

Product Details

| Immunogen: | ADAMTS1 (Thr854-Ser967) |
|---------------|--|
| Isotype: | IgG |
| Specificity: | The antibody is a rabbit polyclonal antibody raised against ADAMTS1. It has been selected for its ability to recognize ADAMTS1 in immunohistochemical staining and western blotting. |
| Purification: | Antigen-specific affinity chromatography |

Target Details

| Target: | ADAMTS1 |
|-------------------|--|
| Alternative Name: | A Disintegrin And Metalloproteinase With Thrombospondin 1 (ADAMTS1) (ADAMTS1 Products) |
| Background: | Alternative Names: C3-C5, METH1 |

| Pathways: |
|-----------|
|-----------|

SARS-CoV-2 Protein Interactome

Application Details

Application Notes:

Western blotting: 1:50-400 Immunocytochemistry in formalin fixed cells: 1:50-500
 Immunohistochemistry in formalin fixed frozen section: 1:50-500 Immunohistochemistry in paraffin section: 1:10-100 Enzyme-linked Immunosorbent Assay: 1:100-1:5000 Optimal working dilutions must be determined by end user.

Comment:


The thermal stability is described by the loss rate. The loss rate was determined by accelerated thermal degradation test, that is, incubate the protein at 37°C for 48h, and no obvious degradation and precipitation were observed. The loss rate is less than 5% within the expiration date under appropriate storage condition.

Restrictions:

For Research Use only


Handling

| Format: | Liquid |
|--------------------|---|
| Concentration: | Lot specific |
| Buffer: | PBS, pH 7.4, containing 0.02 % Sodium azide, 50 % glycerol. |
| Preservative: | Sodium azide |
| Precaution of Use: | WARNING: Reagents contain sodium azide. Sodium azide is very toxic if ingested or inhaled. |
| | Avoid contact with skin, eyes, or clothing. Wear eye or face protection when handling. If skin or |
| | eye contact occurs, wash with copious amounts of water. If ingested or inhaled, contact a |
| | physician immediately. Sodium azide yields toxic hydrazoic acid under acidic conditions. Dilute |
| | azide-containing compounds in running water before discarding to avoid accumulation of |
| | potentially explosive deposits in lead or copper plumbing. |
| Handling Advice: | Avoid repeated freeze-thaw cycles. |
| Storage: | 4 °C |
| Storage Comment: | Store at 2-8 °C for one month. Aliquot and store at -80 °C for 12 months. |
| Expiry Date: | 12 months |




Western Blotting

Image 1.



Immunohistochemistry

Image 2. Used in DAB staining on fromalin fixed paraffinembedded adrenal gland tissue



Immunohistochemistry

Image 3. DAB staining on IHC-P; Samples: Rat Adrenal gland Tissue