antibodies - online.com







anti-APOC1 antibody (AA 34-88)





_						
0	V	0	r٧	/[Θ	M

Quantity:	100 μL	
Target:	APOC1	
Binding Specificity:	AA 34-88	
Reactivity:	Rat	
Host:	Rabbit	
Clonality:	Polyclonal	
Conjugate:	This APOC1 antibody is un-conjugated	
Application:	Western Blotting (WB), Immunohistochemistry (IHC), Immunocytochemistry (ICC)	

Product Details

Immunogen:	APOC1 (Met34-Ala88)
Isotype:	IgG
Specificity:	The antibody is a rabbit polyclonal antibody raised against APOC1. It has been selected for its ability to recognize APOC1 in immunohistochemical staining and western blotting.
Purification:	Antigen-specific affinity chromatography

Target Details

Target:	APOC1	
Alternative Name:	Apolipoprotein C1 (APOC1) (APOC1 Products)	
Background:	Alternative Names: Apo-C1, Truncated apolipoprotein C-I	

Application Details

Application Notes:

Western blotting: 1:50-400 Immunocytochemistry in formalin fixed cells: 1:50-500
 Immunohistochemistry in formalin fixed frozen section: 1:50-500 Immunohistochemistry in paraffin section: 1:10-100 Enzyme-linked Immunosorbent Assay: 1:100-1:5000 Optimal working dilutions must be determined by end user.

Comment:


The thermal stability is described by the loss rate. The loss rate was determined by accelerated thermal degradation test, that is, incubate the protein at 37°C for 48h, and no obvious degradation and precipitation were observed. The loss rate is less than 5% within the expiration date under appropriate storage condition.

Restrictions:

For Research Use only


Handling

Format:	Liquid	
Concentration:	Lot specific	
Buffer:	PBS, pH 7.4, containing 0.02 % Sodium azide, 50 % glycerol.	
Preservative:	Sodium azide	
Precaution of Use:	WARNING: Reagents contain sodium azide. Sodium azide is very toxic if ingested or inhaled.	
	Avoid contact with skin, eyes, or clothing. Wear eye or face protection when handling. If skin or	
	eye contact occurs, wash with copious amounts of water. If ingested or inhaled, contact a	
	physician immediately. Sodium azide yields toxic hydrazoic acid under acidic conditions. Dilute	
	azide-containing compounds in running water before discarding to avoid accumulation of	
	potentially explosive deposits in lead or copper plumbing.	
Handling Advice:	Avoid repeated freeze-thaw cycles.	
Storage:	4 °C	
Storage Comment:	Store at 2-8 °C for one month. Aliquot and store at -80 °C for 12 months.	
Expiry Date:	12 months	



Western Blotting

Image 1.



Immunohistochemistry

Image 2. DAB staining on IHC-P; Samples: Rat Adrenal gland Tissue