antibodies -online.com





anti-SERPINA6 antibody (AA 280-405)

3 Images



Go to Product page

\sim					
	1//	⊃r	V/I	Φ\	Λ

100 μL	
SERPINA6	
AA 280-405	
Human	
Rabbit	
Polyclonal	
This SERPINA6 antibody is un-conjugated	
tochemistry (ICC)	
It has been selected for its estern blotting.	
t	

A, Member 6(Alpha-1 Antiproteinase, Antitrypsin)

Application Details

Western blotting: 1:50-400 Immunocytochemistry in formalin fixed cells: 1:50-500 Immunohistochemistry in formalin fixed frozen section: 1:50-500 Immunohistochemistry in paraffin section: 1:10-100 Enzyme-linked Immunosorbent Assay: 1:100-1:5000 Optimal working dilutions must be determined by end user.


The thermal stability is described by the loss rate. The loss rate was determined by accelerated thermal degradation test, that is, incubate the protein at 37°C for 48h, and no obvious degradation and precipitation were observed. The loss rate is less than 5% within the expiration date under appropriate storage condition.

Restrictions: For Research Use only

Handling

Comment:

Format:	Liquid	
Concentration:	Lot specific	
Buffer:	PBS, pH 7.4, containing 0.02 % Sodium azide, 50 % glycerol.	
Preservative:	Sodium azide	
Precaution of Use:	WARNING: Reagents contain sodium azide. Sodium azide is very toxic if ingested or inhaled.	
	Avoid contact with skin, eyes, or clothing. Wear eye or face protection when handling. If skin or	
	eye contact occurs, wash with copious amounts of water. If ingested or inhaled, contact a	
	physician immediately. Sodium azide yields toxic hydrazoic acid under acidic conditions. Dilute	
	azide-containing compounds in running water before discarding to avoid accumulation of	
	potentially explosive deposits in lead or copper plumbing.	
Handling Advice:	Avoid repeated freeze-thaw cycles.	
Storage:	4 °C	
Storage Comment:	Store at 2-8 °C for one month. Aliquot and store at -80 °C for 12 months.	
Expiry Date:	12 months	




Western Blotting

Image 1.

Western Blotting

Image 2. Western Blot; Sample: Human Liver lysate; Primary Ab: 1μg/ml Rabbit Anti-Human CBG Antibody Second Ab: 0.2μg/mL HRP-Linked Caprine Anti-Rabbit IgG Polyclonal Antibody (Catalog: SAA544Rb19)



Immunohistochemistry

Image 3. Figure.DAB staining on IHC-P. Samples: Human Tissue