

Datasheet for ABIN1858333

anti-E-cadherin antibody (AA 374-628)





Go to Product page

Overview

Quantity:	100 μL
Target:	E-cadherin (CDH1)
Binding Specificity:	AA 374-628
Reactivity:	Rat
Host:	Rabbit
Clonality:	Polyclonal
Conjugate:	This E-cadherin antibody is un-conjugated
Application:	Western Blotting (WB), Immunohistochemistry (IHC), Immunoprecipitation (IP), Immunocytochemistry (ICC)

Product Details

Purpose:	Polyclonal Antibody to E-cadherin
Immunogen:	RPA017Ra02Recombinant Ecadherin
Isotype:	IgG
Specificity:	The antibody is a rabbit polyclonal antibody raised against E-cadherin. It has been selected for its ability to recognize E-cadherin in immunohistochemical staining and western blotting.
Cross-Reactivity:	Human, Mouse
Purification:	Antigen-specific affinity chromatography followed by Protein A affinity chromatography


Target Details

Target:	E-cadherin (CDH1)
Alternative Name:	E-cadherin (CDH1 Products)
Background:	CDHE, CD324, CDH1, Arc1, ECAD, CDH-E, E-CAD, LCAM, UVO, Cadherin 1 Type 1, Uvomorulin, E
	Cadherin, Calcium-Dependent Adhesion Protein, Epithelial, Liver Cell Adhesion Molecule
Pathways:	WNT Signaling, Sensory Perception of Sound, Cell-Cell Junction Organization, Tube Formation
Application Details	
Application Notes:	Western blotting: 0.5-3 μg/mL,lmmunohistochemistry: 5-30 μg/mL,lmmunocytochemistry: 5-
	30 μg/mL,Optimal working dilutions must be determined by end user.
Comment:	The thermal stability is described by the loss rate. The loss rate was determined by accelerated
	thermal degradation test, that is, incubate the protein at 37°C for 48h, and no obvious
	degradation and precipitation were observed. The loss rate is less than 5% within the expiration
	date under appropriate storage condition.
Restrictions:	For Research Use only
Handling	
Format:	Liquid
Concentration:	0.5 mg/mL
Buffer:	PBS, pH 7.4, containing 0.02 % Sodium azide, 50 % glycerol.
Preservative:	Sodium azide
Precaution of Use:	WARNING: Reagents contain sodium azide. Sodium azide is very toxic if ingested or inhaled.
	Avoid contact with skin, eyes, or clothing. Wear eye or face protection when handling. If skin or
	eye contact occurs, wash with copious amounts of water. If ingested or inhaled, contact a
	physician immediately. Sodium azide yields toxic hydrazoic acid under acidic conditions. Dilute
	azide-containing compounds in running water before discarding to avoid accumulation of
	potentially explosive deposits in lead or copper plumbing.
Handling Advice:	Avoid repeated freeze-thaw cycles.
Storage:	4 °C,-20 °C
Storage Comment:	Store at 4°C for frequent use. Stored at -20°C in a manual defrost freezer for two year without
	detectable loss of activity. Avoid repeated freeze-thaw cycles.

Expiry Date:


12 months

Images




Western Blotting

Image 1.



Western Blotting

Image 2. Western Blot; Sample: Rat Serum; Primary Ab: 3μg/ml Rabbit Anti-Rat CDHE Antibody Second Ab: 0.2μg/mL HRP-Linked Caprine Anti-Rabbit IgG Polyclonal Antibody (Catalog: SAA544Rb19)



Immunohistochemistry

Image 3. Figure.DAB staining on IHC-P. Samples: Rat Tissue