

Datasheet for ABIN1858451
anti-COL2 antibody (AA 1259-1449) (APC)



[Go to Product page](#)

Overview

Quantity:	100 µL
Target:	COL2
Binding Specificity:	AA 1259-1449
Reactivity:	Rabbit
Host:	Guinea Pig
Clonality:	Polyclonal
Conjugate:	This COL2 antibody is conjugated to APC
Application:	Western Blotting (WB), Immunohistochemistry (IHC), Immunocytochemistry (ICC), Immunoprecipitation (IP)

Product Details

Purpose:	Polyclonal Antibody to Collagen Type II (COL2)
Immunogen:	COL2 (AA 1259-1449)
Sequence:	MGHHHHHHSG SEF- LK SLNNQIESIR SPEGSRKNPA RTCRDLQLCH PEWKSGDYWI DPNQGCTLDA MKVFCNMETG ETCVYPNPAT VPKNWWSSK SQKQHVWFG ETINGGFHFS YGDSSLPPNT ANVQMTFLRL LSTEGSQNIT YHCKNSVAYL DEATGNLKKA LLLQGSNDVE IRAEGNSRFT YTVLQDGCTK HTGKWGKTV
Isotype:	IgG
Specificity:	The antibody is a cavia polyclonal antibody raised against COL2. It has been selected for its ability to recognize COL2 in immunohistochemical staining and western blotting.
Purification:	Antigen-specific affinity chromatography followed by Protein A affinity chromatography

Target Details

Target:	COL2
Alternative Name:	COL2 (COL2 Products)

Application Details

Application Notes:	Western blotting: 0.01-2 µg/mL,Immunohistochemistry: 5-20 µg/mL,Immunocytochemistry: 5-20 µg/mL,Optimal working dilutions must be determined by end user.
Comment:	The thermal stability is described by the loss rate. The loss rate was determined by accelerated thermal degradation test, that is, incubate the protein at 37°C for 48h, and no obvious degradation and precipitation were observed. The loss rate is less than 5% within the expiration date under appropriate storage condition.
Restrictions:	For Research Use only

Handling

Format:	Liquid
Concentration:	0.5 mg/mL
Buffer:	0.01M PBS, pH 7.4, containing 0.05 % Proclin-300, 50 % glycerol.
Preservative:	ProClin, Sodium azide
Precaution of Use:	WARNING: Reagents contain sodium azide. Sodium azide is very toxic if ingested or inhaled. Avoid contact with skin, eyes, or clothing. Wear eye or face protection when handling. If skin or eye contact occurs, wash with copious amounts of water. If ingested or inhaled, contact a physician immediately. Sodium azide yields toxic hydrazoic acid under acidic conditions. Dilute azide-containing compounds in running water before discarding to avoid accumulation of potentially explosive deposits in lead or copper plumbing.
Handling Advice:	Avoid repeated freeze/thaw cycles
Storage:	4 °C,-20 °C
Storage Comment:	Store at 4°C for frequent use. Stored at -20°C in a manual defrost freezer for two year without detectable loss of activity. Avoid repeated freeze-thaw cycles.
Expiry Date:	12 months