# antibodies -online.com





## anti-CPLX2 antibody (AA 1-134)

2 Images



#### Overview

Quantity:	100 μL
Target:	CPLX2
Binding Specificity:	AA 1-134
Reactivity:	Mouse
Host:	Rabbit
Clonality:	Polyclonal
Conjugate:	This CPLX2 antibody is un-conjugated
Application:	Western Blotting (WB), Immunohistochemistry (IHC), Immunocytochemistry (ICC)

#### **Product Details**

Immunogen:	CPLX2 (Met1-Lys134)
Isotype:	IgG
Specificity:	The antibody is a rabbit polyclonal antibody raised against CPLX2. It has been selected for its ability to recognize CPLX2 in immunohistochemical staining and western blotting.
Purification:	Antigen-specific affinity chromatography

### **Target Details**

Target:	CPLX2
Abstract:	CPLX2 Products
Background:	Alternative Names: 921-L, CPX-2, CPX2, Synaphin-1

Target Details	
Pathways:	Synaptic Vesicle Exocytosis
Application Details	
Application Notes:	Western blotting: 1:50-400 Immunocytochemistry in formalin fixed cells: 1:50-500 Immunohistochemistry in formalin fixed frozen section: 1:50-500 Immunohistochemistry in paraffin section: 1:10-100 Enzyme-linked Immunosorbent Assay: 1:100-1:5000 Optimal working dilutions must be determined by end user.
Comment:	The thermal stability is described by the loss rate. The loss rate was determined by accelerated thermal degradation test, that is, incubate the protein at 37&degC for 48h, and no obvious degradation and precipitation were observed. The loss rate is less than 5% within the expiration date under appropriate storage condition.
Restrictions:	For Research Use only
Handling	
Format:	Liquid
Concentration:	Lot specific
Buffer:	PBS, pH 7.4, containing 0.02 % Sodium azide, 50 % glycerol.
Preservative:	Sodium azide
Precaution of Use:	WARNING: Reagents contain sodium azide. Sodium azide is very toxic if ingested or inhaled.  Avoid contact with skin, eyes, or clothing. Wear eye or face protection when handling. If skin or eye contact occurs, wash with copious amounts of water. If ingested or inhaled, contact a physician immediately. Sodium azide yields toxic hydrazoic acid under acidic conditions. Dilute azide-containing compounds in running water before discarding to avoid accumulation of potentially explosive deposits in lead or copper plumbing.

Store at 2-8 °C for one month. Aliquot and store at -80 °C for 12 months.

Avoid repeated freeze-thaw cycles.

4°C


12 months


Handling Advice:

Storage Comment:

Storage:

Expiry Date:





#### **Western Blotting**

Image 1.

#### **Western Blotting**

Image 2. Western Blot; Sample: Rat Cerebrum lysate; Primary Ab: 2μg/ml Rabbit Anti-Mouse CPLX2 Antibody Second Ab: 0.2μg/mL HRP-Linked Caprine Anti-Rabbit IgG Polyclonal Antibody (Catalog: SAA544Rb19)