antibodies .- online.com





anti-Coagulation Factor IX antibody (AA 232-455)

3 Images



Overview

| OVCIVICV | |
|----------------------|--|
| Quantity: | 100 μL |
| Target: | Coagulation Factor IX (F9) |
| Binding Specificity: | AA 232-455 |
| Reactivity: | Human |
| Host: | Rabbit |
| Clonality: | Polyclonal |
| Conjugate: | This Coagulation Factor IX antibody is un-conjugated |
| Application: | Western Blotting (WB), Immunohistochemistry (IHC), Immunocytochemistry (ICC) |
| Product Details | |
| Immunogen: | F9 (Asp232-Lys455) |
| Isotype: | IgG |
| Specificity: | The antibody is a rabbit polyclonal antibody raised against F9. It has been selected for its ability |
| | to recognize F9 in immunohistochemical staining and western blotting. |
| Purification: | Antigen-specific affinity chromatography |
| | |
| Target Details | |
| Target: | Coagulation Factor IX (F9) |
| Abstract: | F9 Products |
| Background: | Alternative Names: HEMB, FIX, GLA Domain, PTC, Anti Hemophilic Factor B, Christmas Factor, |

| Dlacma Thromb | onlastic Component | Christmas Die | aaca Hamonhilia R |
|---------------|--------------------|---------------|-------------------|

Application Details

Application Notes:

• Western blotting: 1:50-400 Immunocytochemistry in formalin fixed cells: 1:50-500 Immunohistochemistry in formalin fixed frozen section: 1:50-500 Immunohistochemistry in paraffin section: 1:10-100 Enzyme-linked Immunosorbent Assay: 1:100-1:5000 Optimal working dilutions must be determined by end user.

Comment:


The thermal stability is described by the loss rate. The loss rate was determined by accelerated thermal degradation test, that is, incubate the protein at 37°C for 48h, and no obvious degradation and precipitation were observed. The loss rate is less than 5% within the expiration date under appropriate storage condition.

Restrictions:

For Research Use only

Handling

| Format: | Liquid |
|--------------------|--|
| Concentration: | Lot specific |
| Buffer: | PBS, pH 7.4, containing 0.02 % Sodium azide, 50 % glycerol. |
| Preservative: | Sodium azide |
| Precaution of Use: | WARNING: Reagents contain sodium azide. Sodium azide is very toxic if ingested or inhaled. Avoid contact with skin, eyes, or clothing. Wear eye or face protection when handling. If skin or eye contact occurs, wash with copious amounts of water. If ingested or inhaled, contact a physician immediately. Sodium azide yields toxic hydrazoic acid under acidic conditions. Dilute azide-containing compounds in running water before discarding to avoid accumulation of |
| | potentially explosive deposits in lead or copper plumbing. |
| Handling Advice: | Avoid repeated freeze-thaw cycles. |
| Storage: | 4 °C |
| Storage Comment: | Store at 2-8 °C for one month. Aliquot and store at -80 °C for 12 months. |
| Expiry Date: | 12 months |



Western Blotting

Image 1.

Western Blotting

Image 2. Western Blot; Sample: Human 293T cell lysate; Primary Ab: 1μg/ml Rabbit Anti-Human F9 Antibody Second Ab: 0.2μg/mL HRP-Linked Caprine Anti-Rabbit lgG Polyclonal Antibody (Catalog: SAA544Rb19)

Western Blotting

Image 3. Western Blot; Sample: Human HepG2 cell lysate; Primary Ab: 1μg/ml Rabbit Anti-Human F9 Antibody Second Ab: 0.2μg/mL HRP-Linked Caprine Anti-Rabbit lgG Polyclonal Antibody (Catalog: SAA544Rb19)