

Datasheet for ABIN1858803

anti-Fatty Acid Synthase antibody (AA 2257-2504)[Go to Product page](#)**3** Images

Overview

Quantity:	100 µL
Target:	Fatty Acid Synthase (FASN)
Binding Specificity:	AA 2257-2504
Reactivity:	Mouse
Host:	Rabbit
Clonality:	Polyclonal
Conjugate:	This Fatty Acid Synthase antibody is un-conjugated
Application:	Western Blotting (WB), Immunohistochemistry (IHC), Immunoprecipitation (IP), Immunocytochemistry (ICC)

Product Details

Purpose:	Polyclonal Antibody to Fatty Acid Synthase (FASN)
Immunogen:	RPC470Mu01Recombinant Fatty Acid Synthase (FASN)
Isotype:	IgG
Specificity:	The antibody is a rabbit polyclonal antibody raised against FASN. It has been selected for its ability to recognize FASN in immunohistochemical staining and western blotting.
Cross-Reactivity:	Human
Purification:	Antigen-specific affinity chromatography followed by Protein A affinity chromatography

Target Details


Target:	Fatty Acid Synthase (FASN)
Alternative Name:	FASN (FASN Products)
Background:	FAS, OA519, SDR27X1, Short Chain Dehydrogenase/Reductase Family 27X, Member 1, S-acetyltransferase, 3-oxoacyl synthase, Oleoyl hydrolase, Enoyl-acyl-carrier-protein reductase
Pathways:	AMPK Signaling

Application Details

Application Notes:	Western blotting: 0.01-2 µg/mL, Immunohistochemistry: 5-30 µg/mL, Immunocytochemistry: 5-30 µg/mL, Optimal working dilutions must be determined by end user.
Comment:	The thermal stability is described by the loss rate. The loss rate was determined by accelerated thermal degradation test, that is, incubate the protein at 37°C for 48h, and no obvious degradation and precipitation were observed. The loss rate is less than 5% within the expiration date under appropriate storage condition.
Restrictions:	For Research Use only


Handling

Format:	Liquid
Concentration:	0.5 mg/mL
Buffer:	PBS, pH 7.4, containing 0.02 % Sodium azide, 50 % glycerol.
Preservative:	Sodium azide
Precaution of Use:	WARNING: Reagents contain sodium azide. Sodium azide is very toxic if ingested or inhaled. Avoid contact with skin, eyes, or clothing. Wear eye or face protection when handling. If skin or eye contact occurs, wash with copious amounts of water. If ingested or inhaled, contact a physician immediately. Sodium azide yields toxic hydrazoic acid under acidic conditions. Dilute azide-containing compounds in running water before discarding to avoid accumulation of potentially explosive deposits in lead or copper plumbing.
Handling Advice:	Avoid repeated freeze-thaw cycles.
Storage:	4 °C, -20 °C
Storage Comment:	Store at 4°C for frequent use. Stored at -20°C in a manual defrost freezer for two year without detectable loss of activity. Avoid repeated freeze-thaw cycles.




Western Blotting

Image 1.



Immunohistochemistry

Image 2. Used in DAB staining on formalin fixed paraffin-embedded testis tissue



Immunohistochemistry

Image 3. DAB staining on IHC-P; Samples: Mouse Testis Tissue