
antibodies -online.com





anti-Hemopexin antibody (AA 235-378)

2 Images



Go to Product page

Overview

Background:

| Quantity: | 100 μL |
|----------------------|--|
| Target: | Hemopexin (HPX) |
| Binding Specificity: | AA 235-378 |
| Reactivity: | Rat |
| Host: | Rabbit |
| Clonality: | Polyclonal |
| Conjugate: | This Hemopexin antibody is un-conjugated |
| Application: | Western Blotting (WB), Immunohistochemistry (IHC), Immunocytochemistry (ICC) |
| Product Details | |
| Immunogen: | HPX (His235-Leu378) |
| Isotype: | IgG |
| Specificity: | The antibody is a rabbit polyclonal antibody raised against HPX. It has been selected for its ability to recognize HPX in immunohistochemical staining and western blotting. |
| Purification: | Antigen-specific affinity chromatography |
| Target Details | |
| Target: | Hemopexin (HPX) |
| Alternative Name: | Hemopexin (HPX Products) |
| | |

Alternative Names: Haemopexin, Beta-1B-glycoprotein

Target Details

Pathways:

Transition Metal Ion Homeostasis, Regulation of Leukocyte Mediated Immunity, Positive Regulation of Immune Effector Process, Production of Molecular Mediator of Immune Response

Application Details

Western blotting: 1:50-400 Immunocytochemistry in formalin fixed cells: 1:50-500
 Immunohistochemistry in formalin fixed frozen section: 1:50-500 Immunohistochemistry in paraffin section: 1:10-100 Enzyme-linked Immunosorbent Assay: 1:100-1:5000 Optimal working dilutions must be determined by end user.

Comment:


The thermal stability is described by the loss rate. The loss rate was determined by accelerated thermal degradation test, that is, incubate the protein at 37°C for 48h, and no obvious degradation and precipitation were observed. The loss rate is less than 5% within the expiration date under appropriate storage condition.

Restrictions:

For Research Use only


Handling

| Format: | Liquid |
|--------------------|--|
| Concentration: | Lot specific |
| Buffer: | PBS, pH 7.4, containing 0.02 % Sodium azide, 50 % glycerol. |
| Preservative: | Sodium azide |
| Precaution of Use: | WARNING: Reagents contain sodium azide. Sodium azide is very toxic if ingested or inhaled. Avoid contact with skin, eyes, or clothing. Wear eye or face protection when handling. If skin or eye contact occurs, wash with copious amounts of water. If ingested or inhaled, contact a physician immediately. Sodium azide yields toxic hydrazoic acid under acidic conditions. Dilute azide-containing compounds in running water before discarding to avoid accumulation of potentially explosive deposits in lead or copper plumbing. |
| Handling Advice: | Avoid repeated freeze-thaw cycles. |
| Storage: | 4 °C |
| Storage Comment: | Store at 2-8 °C for one month. Aliquot and store at -80 °C for 12 months. |
| Expiry Date: | 12 months |



Western Blotting

Image 1.



Immunohistochemistry

Image 2. Figure.DAB staining on IHC-P. Samples: Rat Tissue