antibodies -online.com





anti-IGLL1 antibody (AA 112-206)






	oo to i roddet page


Overview	
Quantity:	100 μL
Target:	IGLL1
Binding Specificity:	AA 112-206
Reactivity:	Mouse
Host:	Rabbit
Clonality:	Polyclonal
Conjugate:	This IGLL1 antibody is un-conjugated
Application:	Western Blotting (WB), Immunohistochemistry (IHC), Immunocytochemistry (ICC)
Product Details	
Immunogen:	lglL1 (Val112-Ala206)
Isotype:	IgG
Specificity:	The antibody is a rabbit polyclonal antibody raised against IgIL1. It has been selected for its ability to recognize IgIL1 in immunohistochemical staining and western blotting.
Purification:	Antigen-specific affinity chromatography
Target Details	
Target:	IGLL1
Alternative Name:	Immunoglobulin Lambda Like Polypeptide 1 (IgIL1) (IGLL1 Products)
Background:	Alternative Names: CD179b, IGL1, IGL5, IGLL, IGO, IGVPB, VPREB2, CD179 antigen-like family

	member B, Ig lambda-5, Immunoglobulin omega polypeptide, Immunoglobulin-related protein 14.1
Pathways:	Regulation of Cell Size
Application Details	
Application Notes:	Western blotting: 1:50-400 Immunocytochemistry in formalin fixed cells: 1:50-500 Immunohistochemistry in formalin fixed frozen section: 1:50-500 Immunohistochemistry in paraffin section: 1:10-100 Enzyme-linked Immunosorbent Assay: 1:100-1:5000 Optimal working dilutions must be determined by end user.
Comment:	The thermal stability is described by the loss rate. The loss rate was determined by accelerated thermal degradation test, that is, incubate the protein at 37°C for 48h, and no obvious degradation and precipitation were observed. The loss rate is less than 5% within the expiration date under appropriate storage condition.
Restrictions:	For Research Use only
Handling	
Format:	
TOTTIAL.	Liquid
Concentration:	Liquid Lot specific
	<u> </u>
Concentration:	Lot specific
Concentration: Buffer:	Lot specific PBS, pH 7.4, containing 0.02 % Sodium azide, 50 % glycerol.
Concentration: Buffer: Preservative:	Lot specific PBS, pH 7.4, containing 0.02 % Sodium azide, 50 % glycerol. Sodium azide WARNING: Reagents contain sodium azide. Sodium azide is very toxic if ingested or inhaled. Avoid contact with skin, eyes, or clothing. Wear eye or face protection when handling. If skin or eye contact occurs, wash with copious amounts of water. If ingested or inhaled, contact a physician immediately. Sodium azide yields toxic hydrazoic acid under acidic conditions. Dilute azide-containing compounds in running water before discarding to avoid accumulation of
Concentration: Buffer: Preservative: Precaution of Use:	Lot specific PBS, pH 7.4, containing 0.02 % Sodium azide, 50 % glycerol. Sodium azide WARNING: Reagents contain sodium azide. Sodium azide is very toxic if ingested or inhaled. Avoid contact with skin, eyes, or clothing. Wear eye or face protection when handling. If skin or eye contact occurs, wash with copious amounts of water. If ingested or inhaled, contact a physician immediately. Sodium azide yields toxic hydrazoic acid under acidic conditions. Dilute azide-containing compounds in running water before discarding to avoid accumulation of potentially explosive deposits in lead or copper plumbing.
Concentration: Buffer: Preservative: Precaution of Use: Handling Advice:	Lot specific PBS, pH 7.4, containing 0.02 % Sodium azide, 50 % glycerol. Sodium azide WARNING: Reagents contain sodium azide. Sodium azide is very toxic if ingested or inhaled. Avoid contact with skin, eyes, or clothing. Wear eye or face protection when handling. If skin or eye contact occurs, wash with copious amounts of water. If ingested or inhaled, contact a physician immediately. Sodium azide yields toxic hydrazoic acid under acidic conditions. Dilute azide-containing compounds in running water before discarding to avoid accumulation of potentially explosive deposits in lead or copper plumbing. Avoid repeated freeze-thaw cycles.



Western Blotting

Image 1.



Immunohistochemistry

Image 2. Used in DAB staining on fromalin fixed paraffinembedded Liver tissue