antibodies -online.com





anti-IL13 Receptor alpha 1 antibody (AA 201-360)






O	4	D		page
(-()	10	Pron	ויאווו	nane

\sim	
()\/⊝	view
\circ	V I C V V

Quantity:	100 μL	
Target:	IL13 Receptor alpha 1 (IL13RA1)	
Binding Specificity:	AA 201-360	
Reactivity:	Mouse	
Host:	Rabbit	
Clonality:	Polyclonal	
Conjugate:	This IL13 Receptor alpha 1 antibody is un-conjugated	
Application:	Western Blotting (WB), Immunohistochemistry (IHC), Immunocytochemistry (ICC)	
Product Details		
Immunogen:	IL13Ra1 (Ile201-Leu360)	
Immunogen: Isotype:	IL13Ra1 (Ile201-Leu360) IgG	
Isotype:	IgG The antibody is a rabbit polyclonal antibody raised against IL13Ra1. It has been selected for its	
Isotype: Specificity:	IgG The antibody is a rabbit polyclonal antibody raised against IL13Ra1. It has been selected for its ability to recognize IL13Ra1 in immunohistochemical staining and western blotting.	
Isotype: Specificity: Purification:	IgG The antibody is a rabbit polyclonal antibody raised against IL13Ra1. It has been selected for its ability to recognize IL13Ra1 in immunohistochemical staining and western blotting.	
Isotype: Specificity: Purification: Target Details	IgG The antibody is a rabbit polyclonal antibody raised against IL13Ra1. It has been selected for its ability to recognize IL13Ra1 in immunohistochemical staining and western blotting. Antigen-specific affinity chromatography	
Isotype: Specificity: Purification: Target Details Target:	IgG The antibody is a rabbit polyclonal antibody raised against IL13Ra1. It has been selected for its ability to recognize IL13Ra1 in immunohistochemical staining and western blotting. Antigen-specific affinity chromatography IL13 Receptor alpha 1 (IL13RA1)	


Target Details

	Cancer/testis antigen 19	
Pathways:	JAK-STAT Signaling, Positive Regulation of Immune Effector Process, Production of Molecular Mediator of Immune Response	
Application Details		
Application Notes:	Western blotting: 1:50-400 Immunocytochemistry in formalin fixed cells: 1:50-500 Immunohistochemistry in formalin fixed frozen section: 1:50-500 Immunohistochemistry in paraffin section: 1:10-100 Enzyme-linked Immunosorbent Assay: 1:100-1:5000 Optimal working dilutions must be determined by end user.	
Comment:	The thermal stability is described by the loss rate. The loss rate was determined by accelerated thermal degradation test, that is, incubate the protein at 37°C for 48h, and no obvious degradation and precipitation were observed. The loss rate is less than 5% within the expiration date under appropriate storage condition.	
Restrictions:	For Research Use only	
Handling		
Format:	Liquid	
Concentration:	Lot specific	
Buffer:	PBS, pH 7.4, containing 0.02 % Sodium azide, 50 % glycerol.	
Preservative:	Sodium azide	
Precaution of Use:	WARNING: Reagents contain sodium azide. Sodium azide is very toxic if ingested or inhaled. Avoid contact with skin, eyes, or clothing. Wear eye or face protection when handling. If skin or eye contact occurs, wash with copious amounts of water. If ingested or inhaled, contact a physician immediately. Sodium azide yields toxic hydrazoic acid under acidic conditions. Dilute azide-containing compounds in running water before discarding to avoid accumulation of potentially explosive deposits in lead or copper plumbing.	
Handling Advice:	Avoid repeated freeze-thaw cycles.	
Storage:	4°C	
Storage Comment:	Store at 2-8 °C for one month. Aliquot and store at -80 °C for 12 months.	
Expiry Date:	12 months	




Western Blotting

Image 1.



Immunohistochemistry

Image 2. Used in DAB staining on fromalin fixed paraffinembedded testis tissue



Immunohistochemistry

Image 3. DAB staining on IHC-P; Samples: Mouse Liver Tissue