antibodies -online.com





anti-IL18RAP antibody (AA 117-240)

2 Images



Go to	D		
	Pron	ויאווו	mane

\sim			
	IV/E	۱/۱۲	$I \cap V$

Overview	
Quantity:	100 μL
Target:	IL18RAP
Binding Specificity:	AA 117-240
Reactivity:	Mouse
Host:	Rabbit
Clonality:	Polyclonal
Conjugate:	This IL18RAP antibody is un-conjugated
Application:	Western Blotting (WB), Immunohistochemistry (IHC), Immunocytochemistry (ICC)
Product Details	
Product Details Immunogen:	IL18RAP (Gln117-Arg240)
	IL18RAP (Gln117-Arg240) IgG
Immunogen:	
Immunogen: Isotype:	IgG The antibody is a rabbit polyclonal antibody raised against IL18RAP. It has been selected for its
Immunogen: Isotype: Specificity:	IgG The antibody is a rabbit polyclonal antibody raised against IL18RAP. It has been selected for its ability to recognize IL18RAP in immunohistochemical staining and western blotting.
Immunogen: Isotype: Specificity: Purification:	IgG The antibody is a rabbit polyclonal antibody raised against IL18RAP. It has been selected for its ability to recognize IL18RAP in immunohistochemical staining and western blotting.
Immunogen: Isotype: Specificity: Purification: Target Details	IgG The antibody is a rabbit polyclonal antibody raised against IL18RAP. It has been selected for its ability to recognize IL18RAP in immunohistochemical staining and western blotting. Antigen-specific affinity chromatography

like family member B, IL-1R accessory protein-like, Interleukin-1 receptor 7, Interleukin-18 receptor beta

Application Details


Application Notes: Western blotting: 1:50-400 Immunocytochemistry in formalin fixed cells: 1:50-500 Immunohistochemistry in formalin fixed frozen section: 1:50-500 Immunohistochemistry in paraffin section: 1:10-100 Enzyme-linked Immunosorbent Assay: 1:100-1:5000 Optimal working dilutions must be determined by end user. Comment: The thermal stability is described by the loss rate. The loss rate was determined by accelerate.

The thermal stability is described by the loss rate. The loss rate was determined by accelerated thermal degradation test, that is, incubate the protein at 37°C for 48h, and no obvious degradation and precipitation were observed. The loss rate is less than 5% within the expiration date under appropriate storage condition.

Restrictions: For Research Use only


Handling

Format:	Liquid
Concentration:	Lot specific
Buffer:	PBS, pH 7.4, containing 0.02 % Sodium azide, 50 % glycerol.
Preservative:	Sodium azide
Precaution of Use:	WARNING: Reagents contain sodium azide. Sodium azide is very toxic if ingested or inhaled.
	Avoid contact with skin, eyes, or clothing. Wear eye or face protection when handling. If skin or
	eye contact occurs, wash with copious amounts of water. If ingested or inhaled, contact a
	physician immediately. Sodium azide yields toxic hydrazoic acid under acidic conditions. Dilute
	azide-containing compounds in running water before discarding to avoid accumulation of
	potentially explosive deposits in lead or copper plumbing.
Handling Advice:	Avoid repeated freeze-thaw cycles.
Storage:	4 °C
Storage Comment:	Store at 2-8 °C for one month. Aliquot and store at -80 °C for 12 months.
Expiry Date:	12 months



Western Blotting

Image 1.



Immunohistochemistry

Image 2. Used in DAB staining on fromalin fixed paraffinembedded Liver tissue