

Datasheet for ABIN1859721  
**anti-Lactoferrin antibody (AA 64-193)**



[Go to Product page](#)

**3** Images

## Overview

Quantity:	100 µL
Target:	Lactoferrin (LTF)
Binding Specificity:	AA 64-193
Reactivity:	Rat
Host:	Rabbit
Clonality:	Polyclonal
Conjugate:	This Lactoferrin antibody is un-conjugated
Application:	Western Blotting (WB), Immunohistochemistry (IHC), Immunocytochemistry (ICC)

## Product Details

Immunogen:	LTF (Cys64-Gly193)
Isotype:	IgG
Specificity:	The antibody is a rabbit polyclonal antibody raised against LTF. It has been selected for its ability to recognize LTF in immunohistochemical staining and western blotting.
Purification:	Antigen-specific affinity chromatography

## Target Details

Target:	Lactoferrin (LTF)
Alternative Name:	Lactoferrin ( <a href="#">LTF Products</a> )
Background:	Alternative Names: HLF2, LF, GIG12, Lactotransferrin, Growth-inhibiting protein 12,

## Target Details

Talalactoferrin, Lfcin-H, Kaliocin-1

Pathways: [Transition Metal Ion Homeostasis](#)

## Application Details

Application Notes: 

- Western blotting: 1:50-400 Immunocytochemistry in formalin fixed cells: 1:50-500 Immunohistochemistry in formalin fixed frozen section: 1:50-500 Immunohistochemistry in paraffin section: 1:10-100 Enzyme-linked Immunosorbent Assay: 1:100-1:5000 Optimal working dilutions must be determined by end user.

Comment: The thermal stability is described by the loss rate. The loss rate was determined by accelerated thermal degradation test, that is, incubate the protein at 37°C for 48h, and no obvious degradation and precipitation were observed. The loss rate is less than 5% within the expiration date under appropriate storage condition.

Restrictions: For Research Use only

## Handling

Format: Liquid

Concentration: Lot specific

Buffer: PBS, pH 7.4, containing 0.02 % Sodium azide, 50 % glycerol.

Preservative: Sodium azide




Precaution of Use: WARNING: Reagents contain sodium azide. Sodium azide is very toxic if ingested or inhaled. Avoid contact with skin, eyes, or clothing. Wear eye or face protection when handling. If skin or eye contact occurs, wash with copious amounts of water. If ingested or inhaled, contact a physician immediately. Sodium azide yields toxic hydrazoic acid under acidic conditions. Dilute azide-containing compounds in running water before discarding to avoid accumulation of potentially explosive deposits in lead or copper plumbing.

Handling Advice: Avoid repeated freeze-thaw cycles.

Storage: 4 °C

Storage Comment: Store at 2-8 °C for one month. Aliquot and store at -80 °C for 12 months.

Expiry Date: 12 months



### Western Blotting

**Image 1.**

### Western Blotting

**Image 2.** Western Blot; Sample: Rat Serum; Primary Ab: 1µg/ml Rabbit Anti-Rat LTF Antibody Second Ab: 0.2µg/mL HRP-Linked Caprine Anti-Rabbit IgG Polyclonal Antibody (Catalog: SAA544Rb19)

### Immunohistochemistry

**Image 3.** Figure.DAB staining on IHC-P. Samples: Rat Tissue