antibodies -online.com





anti-MANF antibody (AA 28-182)

2 Images



Go to Product page

\sim				
	$ V \cap$	r\/I	19	٨

Uverview	
Quantity:	100 μL
Target:	MANF
Binding Specificity:	AA 28-182
Reactivity:	Human
Host:	Rabbit
Clonality:	Polyclonal
Conjugate:	This MANF antibody is un-conjugated
Application:	Western Blotting (WB), Immunohistochemistry (IHC), Immunocytochemistry (ICC)
Product Details	
Immunogen:	MANF (Gly28-Leu182)

Immunogen:	MANF (Gly28-Leu182)
Isotype:	IgG
Specificity:	The antibody is a rabbit polyclonal antibody raised against MANF. It has been selected for its ability to recognize MANF in immunohistochemical staining and western blotting.
Purification:	Antigen-specific affinity chromatography


Target Details

Target:	MANF
Alternative Name:	Mesencephalic Astrocyte Derived Neurotrophic Factor (MANF) (MANF Products)
Background:	Alternative Names: ARMET, ARP, Arginine Rich Mutated In Early Stage Tumors, Arginine-rich

Expiry Date:


12 months

l arget Details	
	protein
Application Details	
Application Notes:	Western blotting: 1:50-400 Immunocytochemistry in formalin fixed cells: 1:50-500 Immunohistochemistry in formalin fixed frozen section: 1:50-500 Immunohistochemistry in paraffin section: 1:10-100 Enzyme-linked Immunosorbent Assay: 1:100-1:5000 Optimal working dilutions must be determined by end user.
Comment:	The thermal stability is described by the loss rate. The loss rate was determined by accelerated thermal degradation test, that is, incubate the protein at 37°C for 48h, and no obvious degradation and precipitation were observed. The loss rate is less than 5% within the expiration date under appropriate storage condition.
Restrictions:	For Research Use only
Handling	
Format:	Liquid
Concentration:	Lot specific
Concentration: Buffer:	Lot specific PBS, pH 7.4, containing 0.02 % Sodium azide, 50 % glycerol.
Buffer:	PBS, pH 7.4, containing 0.02 % Sodium azide, 50 % glycerol.
Buffer: Preservative:	PBS, pH 7.4, containing 0.02 % Sodium azide, 50 % glycerol. Sodium azide WARNING: Reagents contain sodium azide. Sodium azide is very toxic if ingested or inhaled. Avoid contact with skin, eyes, or clothing. Wear eye or face protection when handling. If skin or eye contact occurs, wash with copious amounts of water. If ingested or inhaled, contact a physician immediately. Sodium azide yields toxic hydrazoic acid under acidic conditions. Dilute azide-containing compounds in running water before discarding to avoid accumulation of
Buffer: Preservative: Precaution of Use:	PBS, pH 7.4, containing 0.02 % Sodium azide, 50 % glycerol. Sodium azide WARNING: Reagents contain sodium azide. Sodium azide is very toxic if ingested or inhaled. Avoid contact with skin, eyes, or clothing. Wear eye or face protection when handling. If skin or eye contact occurs, wash with copious amounts of water. If ingested or inhaled, contact a physician immediately. Sodium azide yields toxic hydrazoic acid under acidic conditions. Dilute azide-containing compounds in running water before discarding to avoid accumulation of potentially explosive deposits in lead or copper plumbing.



Western Blotting

Image 1.



Immunohistochemistry

Image 2. Used in DAB staining on fromalin fixed paraffinembedded Bowels tissue